

# **EXHIBIT A**

US006355432B1

(12) **United States Patent**  
Fodor et al.

(10) Patent No.: **US 6,355,432 B1**  
(45) Date of Patent: **Mar. 12, 2002**

(54) **PRODUCTS FOR DETECTING NUCLEIC ACIDS**

(75) Inventors: **Stephen P. A. Fodor, Palo Alto;**  
**Dennis W. Solas, San Francisco;**  
**William J. Dower, Menlo Park, all of**  
**CA (US)**

(73) Assignee: **Affymetrix Inc., Santa Clara, CA (US)**

(\*) Notice: Subject to any disclaimer, the term of this patent is extended or adjusted under 35 U.S.C. 154(b) by 0 days.

(21) Appl. No.: **09/585,659**

(22) Filed: **Jun. 2, 2000**

**Related U.S. Application Data**

(63) Continuation of application No. 09/362,089, filed on Jul. 28, 1999, which is a division of application No. 09/056,927, filed on Apr. 8, 1998, now Pat. No. 6,197,506, which is a continuation of application No. 08/670,118, filed on Jun. 25, 1996, now Pat. No. 5,800,992, which is a division of application No. 08/168,904, filed on Dec. 15, 1993, now abandoned, which is a continuation of application No. 07/624,114, filed on Dec. 6, 1990, now abandoned, which is a continuation-in-part of application No. 07/492,462, filed on Mar. 7, 1990, now Pat. No. 5,143,854, which is a continuation-in-part of application No. 07/362,901, filed on Jun. 7, 1989, now abandoned.

(51) Int. Cl.<sup>7</sup> ..... **C12Q 1/68; G01N 33/53;**  
**C12M 1/34; C07H 21/04; A61K 38/00**  
(52) U.S. Cl. .... **435/6; 435/7.1; 435/287.2;**  
**435/288.3; 530/300; 530/350; 530/387.1;**  
**536/23.1; 536/24.3**  
(58) Field of Search ..... **536/23.1, 24.3;**  
**435/6, 287.2, 288.3, 7.1; 530/300, 350,**  
**387.1**

(56) **References Cited**

**U.S. PATENT DOCUMENTS**

3,730,844 A 5/1973 Gilham et al. .... 195/103.5 R  
3,849,137 A 11/1974 Barzynski et al. .... 96/97  
3,862,056 A 1/1975 Hartman ..... 252/511  
3,939,350 A 2/1976 Kronick et al. .... 250/365  
4,072,576 A 2/1978 Arwin et al. .... 195/103.5 R  
4,121,222 A 10/1978 Diebold et al. .... 347/7  
4,180,739 A 12/1979 Abu-Shumays ..... 250/461 R  
4,216,245 A 8/1980 Johnson ..... 427/2.13  
4,238,757 A 12/1980 Schenck ..... 357/25  
4,269,933 A 5/1981 Pazos ..... 430/291  
4,314,821 A 2/1982 Rice ..... 23/230 B  
4,327,073 A 4/1982 Huang ..... 424/1  
4,339,528 A 7/1982 Goldman ..... 430/323  
4,342,905 A 8/1982 Fujii et al. .... 250/201  
4,373,071 A 2/1983 Itakura ..... 525/375

(List continued on next page.)

**FOREIGN PATENT DOCUMENTS**

CA 1284931 6/1991  
DE 2242394 3/1974  
DE 3440141 5/1986  
DE 3505287 3/1988  
EP 046 083 2/1982  
EP 088 636 9/1983

EP 103 197 3/1984  
EP 127 438 12/1984  
EP 063 810 3/1986  
EP 174 879 3/1986  
EP 194 132 9/1986  
EP 228 075 7/1987  
EP 245 662 11/1987  
EP 268 237 5/1988  
EP 130 523 6/1988

(List continued on next page.)

**OTHER PUBLICATIONS**

Brenner et al., "In vitro cloning of complex mixtures of DNA on microbeads: Physico-chemical separation of differentially expressed cDNAs", *PNAS*, 02/2000, 97:665-1670.  
Brenner et al., "Gene expression analysis by massively parallel signature sequencing (MPSS) on microbead arrays", *Nature Biotechnol.*, 06/2000, 18:630-634.  
Tyagi, "Taking a census of mRNA populations with microbeads", *Nature Biotechnol.* 06/2000, 18:597-598.  
Miller et al., "Detection of bacteria by hybridization of rRNA with DNA-latex and immunodetection of hybrids" *J Clin Microbiol* 1988, 26:1271-1276.  
Sequencing by Hybridization Workshop, listing of participants and workshop presentation summaries, from workshop held 11/19-20/91.  
"A Sequencing Reality Check," *Science*, 242:1245 (1988).  
"Affymax raises \$25 million to develop high-speed drug discovery system," *Biotechnology News*, 10(3):7-8 (1990).  
"Preparation of fluorescent-labeled DNA and its use as a probe in molecular hybridization," *Bioorg Khim.*, 12(11):1508-1513 (1986).  
Abbott et al., "Manipulation of the Wettability of Surfaces on the 0.1- to 1 -Micrometer Scale Through Micromachining and Molecular Self-Assembly," *Science*, 257:1380-1382 (1992).  
Adams et al., "Complementary DNA Sequencing: Expressed Sequence Tags and Human Genome Project," *Science*, 252(5013):1651-1656 (1991).  
Adams et al., "Photolabile Chelators That "Cage" Calcium with Improved Speed of Release and Pre-Photolysis Affinity," *J. Gen. Physiol.*, p. 9a (12/86).  
Adams et al., "Biologically Useful Chelators That Take Up Ca<sup>2+</sup> upon Illumination," *J. Am. Chem. Soc.*, 111:7957-7968 (1989).  
Ajayaghosh et al., "Solid-Phase Synthesis of N-Methyl- and N-Ethylamides of Peptides Using Photolytically Detachable ((3-Nitro-4((alkylamino)methyl)benzamido)methyl)polystyrene Resin," *J. Org. Chem.*, 55(9):2826-2829 (1990).

(List continued on next page.)

Primary Examiner—Stephanie W. Zilmer  
(74) Attorney, Agent, or Firm—Morgan, Lewis & Bockius LLP

(57) **ABSTRACT**

The present invention provides methods and apparatus for sequencing, fingerprinting and mapping biological macromolecules, typically biological polymers. The methods make use of a plurality of sequence specific recognition reagents which can also be used for classification of biological samples, and to characterize their sources.

22 Claims, 2 Drawing Sheets

## US 6,355,432 B1

Page 2

## U.S. PATENT DOCUMENTS

|             |         |                    |            |             |         |                        |           |
|-------------|---------|--------------------|------------|-------------|---------|------------------------|-----------|
| 4,395,486 A | 7/1983  | Wilson et al.      | 435/6      | 4,865,990 A | 9/1989  | Stead et al.           | 435/803   |
| 4,405,771 A | 9/1983  | Jagur              | 528/266    | 4,868,103 A | 9/1989  | Stavrianopoulos et al. | 435/5     |
| 4,444,878 A | 4/1984  | Paulus             | 435/7      | 4,874,500 A | 10/1989 | Madou et al.           | 204/412   |
| 4,444,892 A | 4/1984  | Malnros            | 436/528    | 4,877,745 A | 10/1989 | Hayes et al.           | 436/166   |
| 4,448,534 A | 5/1984  | Wertz et al.       | 356/435    | 4,886,741 A | 12/1989 | Schwartz               | 435/5     |
| 4,458,066 A | 7/1984  | Caruthers et al.   | 536/27     | 4,888,278 A | 12/1989 | Singer et al.          | 435/6     |
| 4,483,920 A | 11/1984 | Gillespie et al.   | 435/6      | 4,921,805 A | 5/1990  | Gebeyehu et al.        | 435/270   |
| 4,500,707 A | 2/1985  | Caruthers et al.   | 536/27     | 4,923,901 A | 5/1990  | Koester et al.         | 521/53    |
| 4,500,919 A | 2/1985  | Schreiber          | 358/78     | 4,925,785 A | 5/1990  | Wang et al.            | 435/6     |
| 4,516,833 A | 5/1985  | Fusek              | 350/162.12 | 4,931,384 A | 6/1990  | Layton et al.          | 435/7.31  |
| 4,517,338 A | 5/1985  | Urdea et al.       | 525/54.11  | 4,946,942 A | 8/1990  | Fuller et al.          | 530/335   |
| 4,533,682 A | 8/1985  | Tortorello et al.  | 523/414    | 4,965,188 A | 10/1990 | Mullis et al.          | 435/6     |
| 4,537,861 A | 8/1985  | Elings et al.      | 436/518    | 4,973,493 A | 11/1990 | Guire                  | 427/2     |
| 4,542,102 A | 9/1985  | Dattagupta et al.  | 435/6      | 4,979,959 A | 12/1990 | Guire                  | 623/66    |
| 4,555,490 A | 11/1985 | Merril             | 436/86     | 4,981,783 A | 1/1991  | Augenlicht             | 435/6     |
| 4,556,643 A | 12/1985 | Paau et al.        | 435/5      | 4,981,985 A | 1/1991  | Kaplan et al.          | 556/50    |
| 4,562,157 A | 12/1985 | Lowe et al.        | 435/291    | 4,984,100 A | 1/1991  | Takayama et al.        | 360/49    |
| 4,563,419 A | 1/1986  | Ranki et al.       | 435/6      | 4,987,065 A | 1/1991  | Stavrianopoulos et al. | 435/5     |
| 4,569,967 A | 2/1986  | Kornreich et al.   | 525/54.11  | 4,988,617 A | 1/1991  | Landegren et al.       | 435/6     |
| 4,580,895 A | 4/1986  | Patel              | 356/39     | 4,992,383 A | 2/1991  | Farnsworth             | 436/89    |
| 4,584,277 A | 4/1986  | Ullman             | 436/501    | 4,994,373 A | 2/1991  | Stavrianopoulos et al. | 435/6     |
| 4,588,682 A | 5/1986  | Groet et al.       | 435/6      | 5,002,867 A | 3/1991  | Macevitz               | 435/6     |
| 4,591,570 A | 5/1986  | Chang              | 435/7.24   | 5,006,464 A | 4/1991  | Chu et al.             | 435/7.1   |
| 4,598,049 A | 7/1986  | Zelinka et al.     | 422/116    | 5,011,770 A | 4/1991  | Kung et al.            | 435/6     |
| 4,613,566 A | 9/1986  | Potter             | 435/6      | 5,013,669 A | 5/1991  | Peters, Jr. et al.     | 436/518   |
| 4,624,915 A | 11/1986 | Schindler et al.   | 435/4      | 5,021,550 A | 6/1991  | Zieger                 | 530/334   |
| 4,626,684 A | 12/1986 | Landa              | 250/328    | 5,026,773 A | 6/1991  | Steel                  | 525/54.11 |
| 4,631,211 A | 12/1986 | Houghten           | 428/35     | 5,026,840 A | 6/1991  | Dattagupta et al.      | 536/27    |
| 4,637,861 A | 1/1987  | Krull et al.       | 204/1 T    | 5,028,525 A | 7/1991  | Gray et al.            | 435/6     |
| 4,656,127 A | 4/1987  | Mundy              | 435/6      | 5,028,545 A | 7/1991  | Soini                  | 436/501   |
| 4,670,380 A | 6/1987  | Dattagupta         | 435/6      | 5,037,882 A | 8/1991  | Steel                  | 525/54.11 |
| 4,677,054 A | 6/1987  | White et al.       | 435/6      | 5,043,265 A | 8/1991  | Tanke et al.           | 435/6     |
| 4,681,859 A | 7/1987  | Kramer             | 436/501    | 5,047,524 A | 9/1991  | Andrus et al.          | 536/27    |
| 4,683,195 A | 7/1987  | Mullis et al.      | 435/6      | 5,064,754 A | 11/1991 | Mills                  | 435/6     |
| 4,683,202 A | 7/1987  | Mullis             | 435/91     | 5,075,077 A | 12/1991 | Durley                 |           |
| 4,689,405 A | 8/1987  | Frank et al.       | 536/27     | 5,077,085 A | 12/1991 | Schnur et al.          | 427/98    |
| 4,704,353 A | 11/1987 | Humphries et al.   | 435/4      | 5,077,210 A | 12/1991 | Eigler et al.          | 435/176   |
| 4,711,955 A | 12/1987 | Ward et al.        | 536/29     | 5,079,600 A | 1/1992  | Schnur et al.          | 357/4     |
| 4,713,326 A | 12/1987 | Dattagupta et al.  | 435/6      | 5,081,584 A | 1/1992  | Onichinski et al.      | 364/497   |
| 4,713,347 A | 12/1987 | Mitchell et al.    | 436/501    | 5,082,830 A | 1/1992  | Brakel et al.          | 514/44    |
| 4,715,413 A | 12/1987 | Backlund et al.    | 141/94     | 5,091,652 A | 2/1992  | Mathies et al.         | 250/458.1 |
| 4,716,106 A | 12/1987 | Chiswell           | 435/6      | 5,096,807 A | 3/1992  | Leaback                | 435/6     |
| 4,719,179 A | 1/1988  | Barany             | 435/172.1  | 5,100,626 A | 3/1992  | Levin                  | 422/100   |
| 4,719,615 A | 1/1988  | Feyrer et al.      | 369/284    | 5,100,777 A | 3/1992  | Chang                  | 435/7.24  |
| 4,722,906 A | 2/1988  | Guire              | 436/501    | 5,112,962 A | 5/1992  | Letsinger et al.       | 536/27    |
| 4,728,502 A | 3/1988  | Hamill             | 422/116    | 5,141,813 A | 8/1992  | Nelson                 | 428/402   |
| 4,728,591 A | 3/1988  | Clark et al.       | 430/5      | 5,143,854 A | 9/1992  | Pirung et al.          | 436/518   |
| 4,731,325 A | 3/1988  | Palva et al.       | 435/6      | 5,149,625 A | 9/1992  | Church et al.          | 435/6     |
| 4,737,344 A | 4/1988  | Koizumi et al.     | 422/100    | 5,153,319 A | 10/1992 | Caruthers et al.       | 536/27    |
| 4,755,458 A | 7/1988  | Rabbani et al.     | 435/5      | 5,164,319 A | 11/1992 | Hafeman et al.         | 435/287.1 |
| 4,762,881 A | 8/1988  | Kauer              | 525/54.11  | 5,171,695 A | 12/1992 | Ekins                  | 436/501   |
| 4,766,062 A | 8/1988  | Diamond et al.     | 435/6      | 5,188,963 A | 2/1993  | Stapleton              | 435/288.3 |
| 4,767,700 A | 8/1988  | Wallace            | 435/6      | 5,192,980 A | 3/1993  | Dixon et al.           | 356/326   |
| 4,777,019 A | 10/1988 | Dandekar           | 422/68     | 5,200,051 A | 4/1993  | Cozzette et al.        | 204/403   |
| 4,780,504 A | 10/1988 | Buendia et al.     | 525/54.11  | 5,202,231 A | 4/1993  | Drmanac et al.         | 435/6     |
| 4,786,170 A | 11/1988 | Groeblter          | 356/318    | 5,206,137 A | 4/1993  | Ip et al.              | 435/6     |
| 4,786,684 A | 11/1988 | Glass              | 525/54.1   | 5,215,882 A | 6/1993  | Bahl et al.            | 435/6     |
| 4,794,150 A | 12/1988 | Steel              | 525/54.11  | 5,215,889 A | 6/1993  | Schultz                | 435/41    |
| 4,808,508 A | 2/1989  | Platzer            | 430/143    | 5,219,726 A | 6/1993  | Evans                  | 435/6     |
| 4,810,869 A | 3/1989  | Yabe et al.        | 250/201    | 5,225,326 A | 7/1993  | Bresser et al.         | 435/6     |
| 4,811,062 A | 3/1989  | Tabata et al.      | 356/152    | 5,232,829 A | 8/1993  | Longiaru et al.        | 435/6     |
| 4,811,218 A | 3/1989  | Hunkapiller et al. | 204/461    | 5,235,028 A | 8/1993  | Barany et al.          | 528/335   |
| 4,812,512 A | 3/1989  | Buendia et al.     | 525/54.11  | 5,242,974 A | 9/1993  | Holmes                 | 525/54.11 |
| 4,820,630 A | 4/1989  | Taub               | 435/5      | 5,252,743 A | 10/1993 | Barrett et al.         | 548/303.7 |
| 4,822,566 A | 4/1989  | Newman             | 422/68     | 5,256,549 A | 10/1993 | Urdea et al.           | 435/91    |
| 4,833,092 A | 5/1989  | Geysen             | 436/501    | 5,258,506 A | 11/1993 | Urdea et al.           | 536/23.1  |
| 4,844,617 A | 7/1989  | Kelderman et al.   | 356/372    | 5,306,641 A | 4/1994  | Saccocio               | 436/85    |
| 4,846,552 A | 7/1989  | Veldkamp et al.    | 350/162.2  | 5,310,893 A | 5/1994  | Erlich et al.          | 536/24.31 |
| 4,849,513 A | 7/1989  | Smith et al.       | 536/27     | 5,324,633 A | 6/1994  | Fodor et al.           | 435/6     |
| 4,855,225 A | 8/1989  | Fung et al.        | 435/6      | 5,328,824 A | 7/1994  | Ward et al.            | 435/6     |
|             |         |                    |            | 5,348,855 A | 9/1994  | Dattagupta et al.      | 435/6     |

## US 6,355,432 B1

Page 3

|                          |         |                        |           |    |             |         |
|--------------------------|---------|------------------------|-----------|----|-------------|---------|
| 5,384,261 A              | 1/1995  | Winkler et al. ....    | 436/518   | EP | 307 476     | 3/1989  |
| 5,405,783 A              | 4/1995  | Pirung et al. ....     | 436/518   | EP | 319 012     | 6/1989  |
| 5,424,186 A              | 6/1995  | Fodor et al. ....      | 435/6     | EP | 328 256     | 8/1989  |
| 5,424,188 A              | 6/1995  | Schneider et al. ....  | 435/6     | EP | 333 561     | 9/1989  |
| 5,432,099 A              | 7/1995  | Ekins ....             | 436/518   | EP | 337 498     | 10/1989 |
| 5,436,327 A              | 7/1995  | Southern et al. ....   | 536/25.34 | EP | 386 229     | 4/1990  |
| 5,445,934 A              | 8/1995  | Fodor et al. ....      | 435/6     | EP | 373 203     | 6/1990  |
| 5,447,841 A              | 9/1995  | Gray et al. ....       | 435/6     | EP | 392 546     | 10/1990 |
| 5,451,505 A              | 9/1995  | Dollinger              |           | EP | 142 299     | 12/1990 |
| 5,474,796 A              | 12/1995 | Brennan ....           | 427/2.13  | EP | 173 339     | 1/1992  |
| 5,486,452 A              | 1/1996  | Gordon et al. ....     | 435/5     | EP | 171 150     | 3/1992  |
| 5,489,507 A              | 2/1996  | Chehab ....            | 435/6     | EP | 237 362     | 3/1992  |
| 5,489,678 A              | 2/1996  | Fodor et al. ....      | 536/22.1  | EP | 185 547     | 6/1992  |
| 5,492,806 A              | 2/1996  | Drmanac et al. ....    | 435/5     | EP | 260 634     | 6/1992  |
| 5,494,810 A              | 2/1996  | Barany et al. ....     | 435/91.52 | EP | 232 967     | 4/1993  |
| 5,510,270 A              | 4/1996  | Fodor et al. ....      | 436/518   | EP | 235 726     | 5/1993  |
| 5,525,464 A              | 6/1996  | Drmanac et al. ....    | 435/6     | EP | 476 014     | 8/1994  |
| 5,527,681 A              | 6/1996  | Holmes ....            | 435/6     | EP | 225 807     | 10/1994 |
| 5,552,270 A              | 9/1996  | Khrapko et al. ....    | 435/6     | EP | 717 113     | 6/1996  |
| 5,556,961 A              | 9/1996  | Footte et al. ....     | 536/27.1  | EP | 721 016     | 7/1996  |
| 5,561,071 A              | 10/1996 | Hollenberg et al. .... | 437/1     | EP | 535 242     | 9/1997  |
| 5,565,324 A              | 10/1996 | Still                  |           | EP | 848 067     | 6/1998  |
| 5,567,809 A              | 10/1996 | Mandecki               |           | EP | 619 321     | 1/1999  |
| 5,569,584 A              | 10/1996 | Augenlicht ....        | 435/6     | FR | 2559783     | 3/1988  |
| 5,571,639 A              | 11/1996 | Hubbell et al. ....    | 430/5     | GB | 2 129 551   | 5/1984  |
| 5,573,905 A              | 11/1996 | Lerner                 |           | GB | 2156074     | 3/1988  |
| 5,593,839 A              | 1/1997  | Hubbell et al. ....    | 435/6     | GB | 2196476     | 4/1988  |
| 5,599,720 A              | 2/1997  | Ekins ....             | 436/501   | GB | 8810400.5   | 5/1988  |
| 5,604,097 A              | 2/1997  | Brenner                |           | GB | 2233654     | 1/1991  |
| 5,604,099 A              | 2/1997  | Erlich et al. ....     | 435/6     | GB | 2248840     | 9/1992  |
| 5,635,400 A              | 6/1997  | Brenner                |           | JP | 49-110601   | 10/1974 |
| 5,641,634 A              | 6/1997  | Mandecki               |           | JP | 60-248669   | 12/1985 |
| 5,643,728 A              | 7/1997  | Slater et al. ....     | 435/6     | JP | 63-084499   | 4/1988  |
| 5,653,939 A              | 8/1997  | Hollis et al. ....     | 422/50    | JP | 63-223557   | 9/1988  |
| 5,654,413 A              | 8/1997  | Brenner                |           | JP | 1-233447    | 9/1989  |
| 5,667,667 A              | 9/1997  | Southern ....          | 205/687   | NO | P 913186    | 8/1991  |
| 5,667,972 A              | 9/1997  | Drmanac et al. ....    | 435/6     | WO | WO 84/03151 | 8/1984  |
| 5,690,894 A              | 11/1997 | Pinkel                 |           | WO | WO 84/03564 | 9/1984  |
| 5,695,940 A              | 12/1997 | Drmanac et al. ....    | 435/6     | WO | WO 85/01051 | 3/1985  |
| 5,698,393 A              | 12/1997 | Macioszek et al. ....  | 435/5     | WO | WO 86/00991 | 2/1986  |
| 5,700,637 A              | 12/1997 | Southern ....          | 435/6     | WO | WO 86/06487 | 11/1986 |
| 5,707,806 A              | 1/1998  | Shuber                 |           | WO | WO 88/01058 | 2/1988  |
| 5,744,305 A              | 4/1998  | Fodor et al. ....      | 435/6     | WO | WO 88/04777 | 6/1988  |
| 5,751,629 A              | 5/1998  | Nova                   |           | WO | WO 89/05616 | 6/1989  |
| 5,770,367 A              | 6/1998  | Southern               |           | WO | WO 89/08834 | 9/1989  |
| 5,776,737 A              | 7/1998  | Dunn ....              | 435/91.1  | WO | WO 89/10977 | 11/1989 |
| 5,777,888 A              | 7/1998  | Rine et al. ....       | 364/496   | WO | WO 89/11548 | 11/1989 |
| 5,800,992 A              | 9/1998  | Fodor et al. ....      | 435/6     | WO | WO 89/12819 | 12/1989 |
| 5,804,563 A              | 9/1998  | Still                  |           | WO | WO 90/00626 | 1/1990  |
| 5,807,522 A              | 9/1998  | Brown                  |           | WO | WO 90/00887 | 2/1990  |
| 5,807,683 A              | 9/1998  | Brenner                |           | WO | WO 90/15070 | 2/1990  |
| 5,830,645 A              | 11/1998 | Pinkel et al. ....     | 435/6     | WO | WO 90/03382 | 4/1990  |
| 5,843,767 A              | 12/1998 | Beattie ....           | 435/287.1 | WO | WO 90/04652 | 5/1990  |
| 5,846,708 A              | 12/1998 | Hollis et al. ....     | 435/6     | WO | WO 90/05789 | 5/1990  |
| 5,846,719 A              | 12/1998 | Brenner                |           | WO | WO 90/07582 | 7/1990  |
| 5,863,722 A              | 1/1999  | Brenner                |           | WO | WO 91/00868 | 1/1991  |
| 5,869,237 A              | 2/1999  | Ward et al. ....       | 435/6     | WO | WO 91/04266 | 4/1991  |
| 5,871,697 A              | 2/1999  | Rothberg et al. ....   | 422/68.1  | WO | WO 91/07087 | 5/1991  |
| 5,972,619 A              | 10/1999 | Drmanac et al. ....    | 435/6     | WO | WO 92/16655 | 1/1992  |
| 6,018,041 A              | 1/2000  | Drmanac et al. ....    | 536/24.3  | WO | WO 92/10092 | 6/1992  |
| 6,023,540 A              | 2/2000  | Walt                   |           | WO | WO 92/10588 | 6/1992  |
| 6,025,136 A              | 2/2000  | Drmanac et al. ....    | 435/6     | WO | WO 93/02992 | 2/1993  |
| 6,040,166 A              | 3/2000  | Erlich et al. ....     | 435/194   | WO | WO 93/09668 | 5/1993  |
| 6,054,270 A              | 4/2000  | Southern               |           | WO | WO 88/01302 | 6/1993  |
| 6,060,240 A              | 5/2000  | Kamb                   |           | WO | WO 93/11262 | 6/1993  |
| FOREIGN PATENT DOCUMENTS |         |                        |           | WO | WO 93/17126 | 9/1993  |
| EP                       | 281 927 | 9/1988                 |           | WO | WO 93/22456 | 11/1993 |
| EP                       | 228 310 | 10/1988                |           | WO | WO 93/22480 | 11/1993 |
| EP                       | 288 310 | 10/1988                |           | WO | WO 95/00530 | 1/1995  |
| EP                       | 304 202 | 2/1989                 |           | WO | WO 95/11995 | 5/1995  |
|                          |         |                        |           | WO | WO 95/33846 | 12/1995 |

## US 6,355,432 B1

Page 4

|    |             |         |
|----|-------------|---------|
| WO | WO 96/23078 | 8/1996  |
| WO | WO 97/10365 | 3/1997  |
| WO | WO 97/17317 | 5/1997  |
| WO | WO 97/19410 | 5/1997  |
| WO | WO 97/27317 | 7/1997  |
| WO | WO 97/29212 | 8/1997  |
| WO | WO 97/31256 | 8/1997  |
| WO | WO 97/45559 | 12/1997 |
| WO | WO 98/03673 | 1/1998  |
| WO | WO 98/31836 | 7/1998  |
| WO | WO 99/60007 | 11/1999 |
| YU | P-570/87    | 4/1987  |
| YU | 18617/87    | 9/1987  |

## OTHER PUBLICATIONS

- Ajayaghosh et al., "Solid-phase synthesis of C-terminal peptide amides using a photoremovable  $\alpha$ -methylphenacylamido anchoring linkage," *Proc. Ind. Natl. Sci. (Chem.Sci.)*, 100(5):389-396 (1988).
- Ajayaghosh et al., "Polymer-supported Solid-phase Synthesis of C-Terminal Peptide N-Methylamides Using a Modified Photoremovable 3-Nitro-4-N-methylaminomethylpolystyrene Support," *Ind.J.Chem.*, 27B:1004-1008 (1988).
- Ajayaghosh et al., "Polymer-Supported Synthesis of Protected Peptide Segments on a Photosensitive o-Nitro ( $\alpha$ -Methyl)Bromobenzyl Resin," *Tetrahedron*, 44(21):6661-6666 (1988).
- Amit et al., "Photosensitive Protecting Groups of Amino Sugars and Their Use in Glycoside Synthesis. 2-Nitrobenzylloxycarbonylamino and 6-Nitroveratryloxycarbonylamino Derivatives," *J.Org.Chem.*, 39(2):192-196 (1974).
- Amit et al., "Photosensitive Protecting Groups—A Review," *Israel J. Chem.*, 12(1-2):103-113 (1974).
- Anand et al., "A 3.5 genome equivalent multi access YAC library: construction, characterisation, screening and storage," *Nuc. Acids Res.*, 18(8):1951-1956 (1990).
- Anderson et al., "Quantitative Filter Hybridisation," chapter 3 from *Nucleic Acid Hybridization a practical approach*, pp. 73-111, Hamès et al., eds., IRL Press (1985).
- Applied Biosystems, Model 431A Peptide Synthesizer User's manual, Sections 2 and 6, (Aug. 15, 1989).
- Anold et al., "A Novel Universal Support for DNA & RNA Synthesis," abstract from *Federation Proceedings*, 43(7):abstract No. 3669 (1984).
- Atherton et al., *Solid Phase Peptide Synthesis: A Practical Approach*, IRL Press, (1989), tbl. of cont., pp. vii-ix.
- Augenlicht et al., "Cloning and Screening of Sequences Expressed in a Mouse Colon Tumor," *Cancer Research*, 42:1088-1093 (1982).
- Augenlicht et al., "Expression of Cloned Sequences in Biopsies of Human Colonic Tissue and in Colonic Carcinoma Cells Induced to Differentiate in Vitro," *Cancer Res.*, 47:6017-6021 (1987).
- Bains, W., "Hybridization Methods for DNA Sequencing," *Genomics*, 11(2):294-301 (1991).
- Bains et al., "A Novel Method for Nucleic Acid Sequence Determination," *J.Theor.Biol.*, 135:303-307 (1988).
- Bains, W., "Alternative Routes Through the Genome," *Bio-technology*, 8:1251-1256 (1988).
- Balachander et al., "Functionalized Siloxy-Anchored Monolayers with Exposed Amino, Azido, Bromo, or Cyano Groups," *Tetrahed. Ltrs.*, 29(44):5593-5594 (1988).
- Baldwin et al., "New Photolabile Phosphate Protecting Groups," *Tetrahed.*, 46(19):6879-6884 (1990).
- Bannwarth et al., "Laboratory Methods, A System for the Simultaneous Chemical synthesis of Different DNA Fragments on Solid Support," *DNA*, 5(5):413-419 (1986).
- Bannwarth, W., "Gene Technology: a Challenge for a Chemist," *Chimia*, 41(9):302-317 (1987).
- Barany, F., "Genetic disease detection and DNA amplification using cloned thermostable ligase," *PNAS*, 88:189-193 (1991).
- Barltrop et al., "Photosensitive Protective Groups," *Chemical Communications*, pp. 822-823 (1966).
- Barinaga, M., "Will 'DNA Chip' Speed Genome Initiative," *Science*, 253:1489 (1985).
- Bart et al., "Microfabricated Electrohydrodynamic Pumps," *Sensors and Actuators*, A21:-A23:193-197 (1990).
- Bartsh et al., "Cloning of mRNA sequences from the human colon: Preliminary characterisation of defined mRNAs in normal and neoplastic tissues," *Br.J.Can.*, 54:791-798 (1986).
- Baum, R., "Fledgling firm targets drug discovery process," *Chem. Eng. News*, p. 10-11 (1990).
- Beltz et al., "Isolation of Multigene Families and Determination of Homologies by Filter Hybridization Methods," *Methods in Enzymology*, 100:266-285 (1983).
- Benschop, Chem. Abstracts 114(26):256643 (1991).
- Bhatia et al., "New Approach To Producing Patterned Biomolecular Assemblies," *J. American Chemical Society*, 114:4432-4433 (1992).
- Biorad Chromatography Electrophoresis Immunochemistry Molecular Biology HPLC catalog M 1987 pp. 182.
- Blawas et al., "Step-and-Repeat Photopatterning of Protein Features Using Caged-Biotin-BSA: Characterization and Resolution," *Langmuir*, 14(15):4243-4250 (1998).
- Blawas, A.S., "Photopatterning of Protein Features using Caged-biotin-Bovine Serum Albumin," dissertation for Ph.D at Duke University in 1998.
- Bos et al., "Amino-acid substitutions at codon 13 of the N-ras oncogene in human acute myeloid leukaemia," *Nature*, 315:726-730 (1985).
- Boyle et al., "Differential distribution of long and short interspersed element sequences in the mouse genome: Chromosome karyotyping by fluorescence in situ hybridization," *PNAS*, 87:7757-7761 (1990).
- Brock et al., "Rapid fluorescence detection of in situ hybridization with biotinylated bovine herpesvirus-1 DNA probes," *J. Veterinary Diagnostic Invest.*, 1:34-38 (1989).
- Burgi et al., "Optimization in Sample Stacking for High-Performance Capillary Electrophoresis," *Anal. Chem.*, 63:2042-2047 (1991).
- Cameron et al., "Photogeneration of Organic Bases from o-Nitrobenzyl-Derived Carbamates," *J. Am. Chem. Soc.*, 113:4303-4313 (1991).
- Carrano et al., "A High-Resolution, Fluorescence-Based, Semiautomated Method for DNA Fingerprinting," *Genomics*, 4:129-136 (1989).
- Caruthers, M.H., "Gene Synthesis Machines: DNA Chemistry and Its Uses," *Science*, 230:281-285 (1985).
- Chatterjee et al., "Inducible Alkylation of DNA Using an Oligonucleotide-Quinone Conjugates," *Am. J. Chem. Soc.*, 112:6397-6399 (1990).
- Chee et al., "Accessing Genetic Information with High-Density DNA Arrays," *Science*, 274:610-614 (1996).
- Chehab et al., "Detection of sickle cell anaemia mutation by colour DNA amplification," *Lancet*, 335:15-17 (1990).



US 6,355,432 B1

Page 5

- Chehab et al., "Detection of specific DNA sequences by fluorescence amplification: A color complementation assay," *PNAS*, 86:9178-9182 (1989).
- Chetverin et al., "Oligonucleotide Arrays: New Concepts and Possibilities," *Biotechnology*, 12:1093-1099 (1994).
- Church et al., "Multiplex DNA sequencing," *Science*, 240:185-188 (1988).
- Church et al., "Genomic sequencing," *PNAS*, 81:1991-1995 (1984).
- Clevite Corp., Piezoelectric Technology, Data for Engineers.
- Corbett et al., "Reaction of Nitroso Aromatics with Glyoxylic Acid. A New Path to Hydroxamic Acids," *J. Org. Chem.*, 45:2834-2839 (1980).
- Coulson et al., "Toward a physical map of the genome of the nematode *Caenorhabditis elegans*," *PNAS*, 83:7821-7825 (1986).
- Craig et al., "Ordering of cosmid clones covering the Herpes simplex virus type 1 (HSV-1) genome: a test case for fingerprinting the hybridization," *Nuc. Acid. Res.*, 18(9):2653-2660 (1990).
- Cummings et al., "Photoactivable Fluorophores. 1. Synthesis and Photoactivation of o-Nitrobenzyl-Quenched Fluorescent Carbamates," *Tetrahedron Letters*, 29(1):65-68 (1988).
- Dattagupta et al., "Rapid identification of Microorganisms by Nucleic Acid Hybridization after Labeling the Test Sample," *Anal. Biochem.*, 177:85-89 (1989).
- Dattagupta et al., "Nucleic Acid Hybridization: a Rapid Method for the Diagnosis of Infectious Diseases," *Perspectives in Antiinfective Therapy*, eds. Jackson et al., pp. 241-247 (1988).
- Dower et al., "The Search for Molecular Diversity (II): Recombinant and Synthetic Randomized Peptide Libraries," *Ann. Rep. Med. Chem.*, 26:271-280 (1991).
- Diggelmann, "Investigating the VLSIPS synthesis process," Sep. 9, 1994.
- Di Mauro et al., "DNA Technology in Chip Construction," *Adv. Mater.*, 5(5):384-386 (1993).
- Drmanac et al., "An Algorithm for the DNA Sequence Generation from k-Tuple Word Contents of the Minimal Number of Random Fragments," *J. Biomol. Struct. Dyn.*, 8(5):1085-1102 (1991).
- Drmanac et al., "Partial Sequencing by Oligo-Hybridization Concept and Applications in Genome Analysis," 1st Int. Conf. Electrophor., Supercomp., Hum. Genome pp. 60-74 (1990).
- Drmanac et al., "Sequencing by Oligonucleotide Hybridization: A Promising Framework in Decoding of the Genome Program?," 1st Int. Conf. Electrophor., Supercomp., Hum. Genome pp. 47-59 (1990).
- Drmanac et al., "Laboratory Methods, Reliable Hybridization: theory of the Method," *Genomics*, 4:114-128 (1989).
- Drmanac et al., "Sequencing of Megabase Plus DNA by Hybridization: Theory of the Method," abstract of presentation given at Cold Spring Harbor Symposium on Genome Mapping and Sequencing, Apr. 27, 1988 thru May 1, 1988.
- Dulcey et al., "Deep UV Photochemistry of Chemisorbed Monolayers: Patterned Coplanar Molecular Assemblies," *Science*, 252:551-554 (1991).
- Duncan et al., "Affinity Chromatography of a Sequence-Specific DNA Binding Protein Using Teflon-Linked Oligonucleotides," *Analytical Biochemistry*, 169:104-108 (1988).
- Effenhauser et al., "Glass Chips for High-speed Capillary Electrophoresis Separations with Submicrometer Plate Heights," *Anal. Chem.*, 65:2637-2642 (1993).
- Effenhauser et al., "High-Speed Separation of Antisense Oligonucleotides on a Micromachined Capillary Electrophoresis Device," *Anal. Chem.*, 66:2949-2953 (1994).
- Ekins et al., "High Specific Activity Chemiluminescent and Fluorescent Markers: their Potential Application to High Sensitivity and 'Multi-analyte' Immunoassays," *J. Bioluminescence Chemiluminescence*, 4:59-78 (1989).
- Ekins et al., "Development of Microspot Multi-Analyte Ratiometric Immunoassay Using dual Fluorescent-Labelled Antibodies," *Anal. Chimica Acta*, 227:73-96 (1989).
- Ekins et al., "Multianalyte Microspot Immunoassay-Microanalytical 'Compact Disk' of the Future," *Clin. Chem.*, 37(11):1955-1967 (1991).
- Ekins, R.P., "Multi-Analyte immunoassay\*," *J. Pharmaceutical Biomedical Analysis*, 7(2):155-168 (1989).
- Ekins et al., "Fluorescence Spectroscopy and its Application to a New Generation of High Sensitivity, Multi-Microspot, Multianalyte, Immunoassay," *Clin. Chim. Acta*, 194:91-114 (1990).
- Elder, J.K., "Analysis of DNA Oligonucleotide Hybridization Data by Maximum Entropy," *Maximum Entropy and Bayesian Methods*, eds. Mohammad-Djafari and Demoment, Kluwer, Dordrecht, pp. 363-371 (1992).
- Ellis, R.W., "The Applications of Synthetic Oligonucleotides to Molecular Biology," *Pharmaceutical Research*, 3(4):195-207 (1986).
- Evans et al., "Microfabrication for Automation of Molecular processes in Human Genome Analysis," *Clin. Chem.*, 41(11):1681 (1995).
- Evans et al., "Physical mapping of complex genomes by cosmid multiplex analysis," *PNAS*, 86:5030-5034 (1989).
- Ezaki et al., "Small-Scale DNA Preparation for Rapid Genetic Identification of *Campylobacter* Species without Radioisotope," *Microbiol. Immunology*, 32(2):141-150 (1988).
- Fan et al., "Mapping small DNA sequences by fluorescence in situ hybridization directly on banded metaphase chromosomes," *PNAS*, 87(16):6223-6227 (1990).
- Fan et al., "Micromachining of Capillary Electrophoresis Injectors and Separators on Glass Chips and Evaluation of Flow at Capillary Intersections," *Anal. Chem.*, 66:177-184 (1994).
- Feinberg et al., Addendum to "A technique for Radiolabeling DNA Restriction Endonuclease Fragments to High Specific Activity," *Anal. Biochem.*, 137:266-267 (1984).
- Fettingner et al., "Stacked modules for micro flow systems in chemical analysis: concept and studies using an enlarged model," *Sensors and Actuators*, B17:19-25 (1993).
- Flanders et al., "A new interferometric alignment technique," *App. Phys. Lett.*, 31(7):426-429 (1977).
- Fodor et al., "Multiplexed biochemical assays with biological chips," *Nature*, 364:555-556 (1993).
- Fodor et al., "Light-directed, Spatially Addressable Parallel Chemical Synthesis," *Science*, 251:767-773 (1991).
- Forman et al., "Thermodynamics of Duplex Formation and Mismatch Discrimination on Photolithographically Synthesized Oligonucleotide Arrays," chapter 13pgs. 206-228 from *Molecular Modeling of Nucleic Acids*, ACS Symposium Series 682, 4/13-17/97, Leontis et al., eds.

US 6,355,432 B1

Page 6

- Frank et al., "Simultaneous Multiple Peptide Synthesis Under Continuous flow Conditions on Cellulose Paper Discs as Segmental Solid Supports," *Tetrahedron*, 44(19):6031-6040 (1988).
- Frank et al., "Automation of DNA Sequencing Reactions and Related Techniques: A Workstation for Micromanipulation of Liquids," *Bio/Technology*, 6:1211-1212 (1988).
- Frank et al., "Simultaneous Synthesis and Biological Applications of DNA Fragments: An Efficient and Complete Methodology," *Methods in Enzymology*, 154:221-250 (1987).
- Fuhr et al., "Travelling wave-driven microfabricated electrohydrodynamic pumps for liquids," *J. Micromech. Microeng.*, 4:217-226 (1994).
- Fuller et al., "Urethane-Protected Amino Acid N-Carboxy Anhydrides and Their Use in Peptide Synthesis," *J. Amer. Chem. Soc.*, 112(20):7414-7416 (1990).
- Furka et al., "General method for rapid synthesis of multi-component peptide mixtures," *Int. J. Peptide Protein Res.*, 37:487-493 (1991).
- Furka et al., "Cornucopia of Peptides by Synthesis," 14th Int. Congress of Biochem. abst.# FR:013, 7/10-15/88 Prague, Czechoslovakia.
- Furka et al., "More Peptides by Less Labour," abst. 288, Int. Symp. Med. Chem., Budapest Hungary.
- Gait, eds., pp. 1-115 from *Oligonucleotide Synthesis: A Practical Approach*, IRL Press, (1984).
- Gazard et al., "Lithographic Technique Using Radiation-Induced Grafting of Acrylic Acid into Poly(Methyl Methacrylate) Films," *Polymer Engineering and Science*, 20(16):1069-1072 (1980).
- Gergen et al., "Filter replicas and permanent collections of recombinant DNA plasmids," *Nuc. Acids Res.*, 7(8):2115-2137 (1979).
- Getzoff et al., "Mechanisms of Antibody Binding to a Protein," *Science*, 235:1191-1196 (1987).
- Geysen et al., "Strategies for epitope analysis using peptide synthesis," *J. Immunol. Meth.*, 102:259-274 (1987).
- Geysen et al., "Use of peptide synthesis to probe viral antigens for epitopes to a resolution of a single amino acid," *PNAS*, 81:3998-4002 (1984).
- Geysen et al., "A synthetic strategy for epitope mapping," from *Peptides: Chem. & Biol.*, Proc. of 10th Am. Peptide Symp., 5/23-28/87, pp. 519-523, (1987).
- Geysen, "Antigen-antibody interactions at the molecular level: adventures in peptide synthesis," *Immunol. Today*, 6(12):364-369 (1985).
- Geysen et al., "Cognitive Features of Continuous Antigenic Determinants," from *Synthetic Peptides: Approaches to Biological Probes*, pp. 19-30, (1989).
- Geysen et al., "Chemistry of Antibody Binding to a Protein," *Science*, 235:1184-1190 (1987).
- Geysen et al., "The delineation of peptides able to mimic assembled epitopes," 1986 CIBA Symp., pp. 130-149.
- Geysen et al., "Cognitive Features of Continuous Antigenic Determinants," *Mol. Recognit.*, 1(1):1-10 (1988).
- Geysen et al., "A Prio Ri Delineation of a Peptide Which Mimics A Discontinuous Antigenic Determinant," *Mol. Immunol.*, 23(7):709-715 (1986).
- Ghosh et al., "Covalent attachment of oligonucleotides to solid supports," *Nuc. Acids Res.*, 15(13):5353-5373 (1987).
- Gilon et al., "Backbone Cyclization: A New Method for Conferring Conformational Constraint on Peptides," *Biopolymers*, 31(6):745-750 (1991).
- Gingeras et al., "Hybridization properties of immobilized nucleic acids," *Nuc. Acids Res.*, 15(13):5373-5390 (1987).
- Gummerlock et al., "RAS Enzyme-Linked Immunoblot Assay Discriminates p21 Species: A Technique to Dissect Gene Family Expression," *Anal. Biochem.*, 180:158-168 (1989).
- Gurney et al., "Activation of a potassium current by rapid photochemically generated step increases of intracellular calcium in rat sympathetic neurons," *PNAS*, 84:3496-3500 (1987).
- Haase et al., "Detection of Two Viral Genomes in Single Cells by Double-Label Hybridization in Situ and Color Microradioautography," *Science*, 227:189-192 (1985).
- Hacia, et al., "Two color hybridization analysis using high density oligonucleotide arrays and energy transfer dyes," *Nuc. Acids Res.*, 26(16):3865-3866 (1998).
- Hack, M.L., "Conics Formed to Make Fluid & Industrial Gas Micromachines," *Genetic Engineering News*, 15(18):1, 29 (1995).
- Hagedorn et al., "Pumping of Water Solutions in Microfabricated Electrohydrodynamic Systems," from *Micro Electro Mechanical Systems conference in Travemunde Germany* (1992).
- Hames et al., *Nuclear acid hybridization, a practical approach*, cover page and table of contents (1985).
- Hanahan et al., "Plasmid Screening at High Colony Density," *Meth. Enzymology*, 100:333-342 (1983).
- Hanahan et al., "Plasmid screening at high colony density," *Gene*, 10:63-67 (1980).
- Haridasan et al., "Peptide Synthesis using Photolytically Cleavable 2-Nitrobenzyloxycarbonyl Protecting Group," *Proc. Indian Natn. Sci. Acad.*, 53A(6):717-728 (1987).
- Harrison et al., "Capillary Electrophoresis and Sample Injection Systems Integrated on a Planar Glass Chip," *Anal. Chem.*, 64:1926-1932 (1992).
- Harrison et al., "Micromachining a Minaturized Capillary Electrophoresis-Based Chemical Analysis System on a Chip," *Science*, 261:895-897 (1993).
- Harrison et al., "Towards minaturized electrophoresis and chemical analysis systems on silicon: an alternative to chemical sensors," *Sensors and Actuators*, B10:107-116 (1993).
- Harrison et al., "Rapid separation of fluorescein derivatives using a micromachined capillary electrophoresis system," *Analytica Chimica Acta*, 283:361-366 (1993).
- Hellberg et al., "Minimum analogue peptide sets (MAPS) for quantitative structure-activity relationships," *Int. J. Peptide Protein Res.*, 37:414-424 (1991).
- Hilser et al., "Protein and peptide mobility in capillary zone electrophoresis, A comparison of existing models and further analysis," *J. Chromatography*, 630:329-336 (1993).
- Ho et al., "Highly Stable Biosensor Using an Artificial Enzyme," *Anal. Chem.*, 59:536-537 (1987).
- Hochgeschwender et al., "Preferential expression of a defined T-cell receptor  $\beta$ -chain gene in hapten-specific cytotoxic T-cell clones," *Nature*, 322:376-378 (1986).
- Hodgson, J., "Assays A La Photolithography," *Biotech.*, 9:419 (1991).
- Hodgson et al., "Hybridization probe size control: optimized 'oligolabelling'," *Nuc. Acids Res.*, 15(15):6295 (1987).
- Hopman et al., "Bi-color detection of two target DNAs by non-radioactive in situ hybridization," *Histochem.*, 85:1-4 (1986).

US 6,355,432 B1

Page 7

- Iwamura et al., "1-Pyrenylmethyl Esters, Photolabile Protecting Groups for Carboxylic Acids," *Tetrahedron Lett.*, 28(6):679-682 (1987).
- Iwamura et al., "1-( $\alpha$ -Diazobenzyl)pyrene: A Reagent for Photolabile and Fluorescent Protection of Carboxyl Groups of Amino Acids and Peptides," *Synlett*, p. 35-36 (1991).
- Jacobson et al., "Effects of Injection Schemes and Column Geometry on the Performance of Microchip Electrophoresis Devices," *Anal. Chem.*, 66:1107-1113 (1994).
- Jacobsen et al., "Open Channel Electrochromatography on a Microchip," *Anal. Chem.*, 66:2369-2373 (1994).
- Jacobson et al., "Microchip Capillary Electrophoresis with an Integrated Postcolumn Reactor" *Anal. Chem.*, 66:3472-3476 (1994).
- Jacobson et al., "Precolumn Reactions with Electrophoretic Analysis Integrated on a Microchip," *Anal. Chem.*, 66:4127-4132 (1994).
- Jacobson et al., "Microfabricated chemical measurement systems," *Nature Medicine*, 1(10):1093-1096 (1995).
- Jacobsen et al., "Fused Quartz Substrates for Microchip Electrophoresis," *Anal. Chem.*, 67:2059-2063 (1995).
- Jacobson et al., "High-Speed Separations on a Microchip," *Anal. Chem.*, 66:1114-1118 (1994).
- Jacobson et al., "Microchip electrophoresis with sample stacking," *Electrophoresis*, 16:481-486 (1995).
- Jayakumari, "Peptide synthesis in a triphasic medium catalysed by papain immobilized on a crosslinked polystyrene support," *Indian J. Chemistry*, 29B:514-517 (1990).
- Jovin et al., "Luminescence Digital Imaging Microscopy," *Ann. Rev. Biophys. Biophys. Chem.*, 18:271-308 (1989).
- Kafatos et al., "Determination of nucleic acid sequence homologies and relative concentrations by a dot hybridization procedure," *Nuc. Acids Res.*, 7(6):1541-1553 (1979).
- Kaiser et al., "Peptide and Protein Synthesis by Segment Synthesis-Condensation," *Science*, 243:187-192 (1989).
- Kaplan et al., "Photolabile chelators for the rapid photorelease of divalent cations," *PNAS*, 85:6571-6575 (1988).
- Karube, "Micro-biosensors based on silicon fabrication technology," chapter 25 from *Biosensors: Fundamentals and Applications*, Turner et al., eds., Oxford Publ., 1987, pp. 471-480 (1987).
- Kates et al., "A Novel, Convenient, Three-dimensional Orthogonal Strategy for Solid-Phase Synthesis of Cyclic Peptides 1-3," *Tetrahed. Letters*, 34(10):1549-1552 (1993).
- Kerkof et al., "A Procedure for Making Simultaneous Determinations of the Relative Levels of Gene Transcripts in Tissues or Cells," *Anal. Biochem.*, 188:349-355 (1990).
- Khrapko et al., "An Oligonucleotide hybridization approach to DNA sequencing," *FEBS Lett.*, 256(1,2):118-122 (1989).
- Khrapko et al., "A method for DNA sequencing by hybridization with oligonucleotide matrix," *DNA Seq. Map.*, 1:375-388 (1991).
- Kidd et al., " $\alpha_1$ -Antitrypsin deficiency detection by direct analysis of the mutation in the gene," *Nature*, 304:230-234 (1983).
- Kievits et al., "Rapid subchromosomal localization of cosmids by nonradioactive in situ hybridization," *Cytogenetics Cell Genetics*, 53(2-3):134-136 (1990).
- Kimura et al., "An Immobilized Enzyme Membrane Fabrication Method using an Ink Jet Nozzle," *Biosensors*, 4:41-52 (1988).
- Kimura et al., "An Integrated SOS/FET Multi-Biosensor," *Sensors & Actuators*, 9:373-387 (1986).
- Kitazawa et al., "In situ DNA-RNA hybridization using in vivo bromodeoxyuridine-labeled DNA probe," *Histochemistry*, 92:195-199 (1989).
- Kleinfeld et al., "Controlled Outgrowth of Dissociated Neurons on Patterned Substrates," *J. Neurosci.*, 8(11):4098-4120 (1988).
- Knight, P., "Materials and Methods/Microsequencers for Proteins and Oligosaccharides," *BioTech.*, 7:1075-76 (1989).
- Kohara et al., "The Physical Map of the Whole *E. coli* Chromosome: Application of a New Strategy for Rapid Analysis and Sorting of a Large Genomic Library," *Cell*, 50:495-508 (1987).
- Krile et al., "Multiplex holography with chirp-modulated binary phase-coded reference-beam masks," *Applied Opt.*, 18(1):52-56 (1979).
- Labat, I., "Subfragments as an informative characteristic of the DNA molecule—computer simulation," research report submitted to the University of Belgrade College of Natural Sciences and Mathematics, (1988).
- Lander et al., "Genomic Mapping by Fingerprinting Random Clones: A Mathematical Analysis," *Genomics*, 2:231-239 (1988).
- Lainer et al., "Human Lymphocyte Subpopulations Identified by Using Three-Color Immunofluorescence and Flow Cytometry Analysis: Correlation of Leu-2, Leu-3, Leu-7, Leu-8, and Leu-11 with Surface Antigen Expression," *Journal of Immunology*, 132(1):151-156 (1984).
- Lam et al., "A new type of synthetic peptide library for identifying ligand-binding activity," *Nature*, 354:82-84 (1991).
- Laskey et al., "Messenger RNA prevalence in sea urchin embryos measured with cloned cDNAs," *PNAS*, 77(9):5317-5321 (1980).
- Lee et al., "synthesis of a Polymer Surface Containing Covalently Attached Triethoxysilane Functionality: Adhesion to Glass," *Macromolecules*, 21:3353-3356 (1988).
- Lehrach et al., "Labelling oligonucleotides to high specific activity (I)," *Nuc. Acids Res.*, 17(12):4605-4610 (89).
- Lehrach et al., "Phage Vectors—EMBL Series," *Meth. Enzymology*, 153:103-115 (1987).
- Lehrach et al., "Hybridization Fingerprinting in Genome Mapping and Sequencing," *Genome Analysis vol. 1: Genetic and Physical Mapping*, Cold Spring Harbor Laboratory Press, pp. 39-81 (1990).
- Levy, M.F., "Preparing Additive Printed Circuits," *IBM Tech. Discl. Bull.*, 9(11):1473 (1967).
- Lewin, Benjamin, eds., *Genes*, third edition, John Wiley & Sons, cover page, preface and table of contents, (1987).
- Lichter et al., "High-Resolution Mapping of Human Chromosome 11 by in Situ hybridization with Cosmid Clones," *Science*, 247:64-69 (1990).
- Lichter et al., "Fluorescence in situ hybridization with Alu and L1 polymerase chain reaction probes for rapid characterization of human chromosomes in hybrid cell lines," *PNAS*, 87:6634-6638 (1990).
- Lichter et al., "Rapid detection of human chromosome 21 aberrations by in situ hybridization," *PNAS*, 85:9664-9668 (1988).
- Lichter et al., "Is non-isotopic in situ hybridization finally coming of age," *Nature*, 345:93-94 (1990).
- Lieberman et al., "A Light source Smaller Than the Optical Wavelength," *Science*, 247:59-61 (1990).



US 6,355,432 B1

Page 8

- Lipshutz et al., "Using Oligonucleotide Probe Arrays To Access Genetic Diversity," *BioTech.*, 19(3):442-7 (1995).
- Little, P., "Clone maps made simple," *Nature*, 346:611-612 (1990).
- Liu et al., "Sequential Injection Analysis in Capillary Format with an Electroosmotic Pump," *Talanta*, 41(11):1903-1910 (1994).
- Lockhart et al., "Expression monitoring by hybridization to high-density oligonucleotide arrays," *Nat. Biotech.* 14:1675-1680 (1996).
- Logue et al., "General Approaches to Mask Design for Binary Optics," *SPIE*, 1052:19-24 (1989).
- Loken et al., "three-color Immunofluorescence Analysis of Leu Antigens on Human Peripheral Blood Using Two Lasers on a Fluorescence-Activated Cell Sorter," *Cytoetry*, 5:151-158 (1984).
- Love et al., "Screening of  $\lambda$  Library for Differentially Expressed Genes Using in Vitro Transcripts," *Anal. Biochem.*, 150:429-441 (1985).
- Lowe, C.R., "Biosensors," *Trends in Biotech.*, 2:59-65 (1984).
- Lowe, C.R., "An Introduction to the Concepts and Technology of Biosensors," *Biosensors*, 1:3-16 (1985).
- Lowe, C. R., *Biotechnology and Crop Improvement and Protection*, BCPC Publications, pp. 131-138 (1986).
- Lowe et al., "Solid-Phase Optoelectronic Biosensors," *Methods in Enzymology*, 137:338-347 (1988).
- Lowe, C.R., "Biosensors," *Phil. Tran. R. Soc. Lond.*, 324:487-496 (1989).
- Lu et al., "Differential screening of murine ascites cDNA libraries by means of in vitro transcripts of cell-cycle-phase-specific cDNA and digital image processing," *Gene*, 86:185-192 (1990).
- Luo, J. et al., "Improving the fidelity of *Thermus thermophilus* DNA ligase," *Nuc.Acids Res.*, 24(14):3071-3078 (1996).
- Lysov et al., "A new method for determining the DNA nucleotide sequence by hybridization with oligonucleotides," *Doklady Biochem.*, 303(1-6):436-438 (1989).
- Lysov et al., "DNA Sequencing by Oligonucleotide Hybridization," First International Conference on Electrophoresis, Supercomputing and the Human Genome, 4/10-13/90 p. 157.
- MacDonald et al., "A Rapid ELISA for Measuring Insulin in a Large Number of Research Samples," *Metabolism*, 38(5):450-452 (1989).
- Mairanovsky, V.G., "Electro-Deprotection-Electrochemical Removal of Protecting Groups\*\*," *Agnew. Chem. Int. Ed. Engl.*, 15(5):281-292 (1976).
- Manz et al., "Miniaturized Total Chemical Analysis Systems: a Novel Concept for Chemical Sensing," *Sensors and Actuators*, B1:244-248 (1990).
- Manz et al., "Micromachining of monocrystalline silicon and glass for chemical analysis systems, A look into next century's technology or just a fashionable craze?," *Trends in Analytical Chem.*, 10(5):144-149 (1991).
- Manz et al., "Planar chips technology for miniaturization and integration of separation techniques into monitoring systems, Capillary electrophoresis on a chip," *J. Chromatography*, 593:253-258 (1992).
- Manz et al., "Planar Chips Technology for Miniaturization of Separation Systems: A Developing Perspective in Chemical Monitoring," chapter 1, 1-64 (1993).
- Manz et al., "Electroosmotic pumping and electrophoretic separations for miniaturized chemical analysis systems," *J. Micromech. Microeng.*, 4:257-265 (1994).
- Masiakowski et al., "Cloning of cDNA sequences of hormone-regulated genes from the MCF-7 human breast cancer cell line," *Nuc. Acids Res.*, 10(24):7895-7903 (1982).
- Matsumoto et al., "Preliminary Investigation of Micropumping Based on Electrical Control of Interfacial Tension," *IEEE*, pp. 105-110 (1990).
- Matsuzawa et al., "Containment and-growth of neuroblastoma cells on chemically patterned substrates," *J. Neurosci. Meth.*, 50:253-260 (1993).
- Matthes et al., "Simultaneous rapid chemical synthesis of over one hundred oligonucleotides on a microscale," *EMBO J.*, 3(4):801-805 (1984).
- McCray et al., "Properties and Uses of Photoreactive Caged Compounds," *Ann. Rev. Biophys. Biophys. Chem.*, 18:239-270 (1989).
- McGall et al., "The Efficiency of Light-Directed Synthesis of DNA Arrays on Glass Substrates," *J. American Chem. Soc.*, 119(22):5081-5090 (1997).
- McGillis, VLSI Technology, Sze, eds., Chapter 7, "Lithography," pp. 267-301 (1983).
- McMurray, J.S., "Solid Phase Synthesis of a Cyclic Peptide Using Fmoc Chemistry," *Tetrahedron Letters*, 32(52):7679-7682 (1991).
- Meinkoth et al., "Review: Hybridization of Nucleic Acids Immobilized on solid Supports," *Analytical Biochem.*, 138:267-284 (1984).
- Melcher et al., "Traveling-Wave Bulk Electroconvection Induced across a Temperature Gradient," *Physics of Fluids*, 10(6):1178-1185 (1967).
- Merrifield, R.B., "Solid Phase peptide Synthesis. I. The Synthesis of a Tetrapeptide," *J. Am. Chem. Soc.*, 85:2149-2154 (1963).
- Michiels et al., "Molecular approaches to genome analysis: a strategy for the construction of ordered overlapping clone libraries," *CABIOS*, 3(3):203-10 (1987).
- Mirzabekov, A.D., "DNA sequencing by hybridization—a megasequencing method and a diagnostic tool?," *TIBTECH*, 12:27-32 (1994).
- Miyada et al., "Oligonucleotide Hybridization Techniques," *Meth. Enzymology*, 154:94-107 (1987).
- Monaco et al., "Human Genome Linking with Cosmids and Yeast Artificial Chromosomes", abstract from CSHS, p. 50, (1989).
- Morita et al., "Direct pattern fabrication on silicone resin by vapor phase electron beam polymerization," *J. Vac. Sci. Technol.*, B1(4):1171-1173 (1983).
- Morrison et al., "Solution-Phase Detection of Polynucleotides Using Interacting Fluorescent Labels and Competitive Hybridization," *Anal. Biochem.*, 183:231-244 (1989).
- Munegumi et al., "thermal Synthesis of Polypeptides from N-Boc-Amino Acid (Aspartic Acid,  $\beta$ -Aminoglutaric Acid) Anhydrides," *Chem. Letters*, pp. 1643-1646 (1988).
- Mutter et al., "Impact of Conformation on the Synthetic Strategies for Peptide Sequences," pp. 217-228 from *Chemistry of Peptides and Proteins*, vol. 1, Proceedings of the Third USSR-FRG Symp., in USSR (1982).
- Nakamori et al., "A Simple and Useful Method for Simultaneous Screening of Elevated Levels of Expression of a Variety of Oncogenes in Malignant Cells," *Jpn. J. Cancer Res.*, 79:1311-1317 (1988).

IAFP00000662

US 6,355,432 B1

Page 9

- Nederlof et al., "Multiple Fluorescence In Situ Hybridization," *Cytometry*, 11:126-131 (1990).
- Nederlof et al., "Three-Color Fluorescence In Situ Hybridization for the Simultaneous Detection of Multiple Nucleic Acids Sequences," *Cytometry*, 10:20-27 (1989).
- Nizetic et al., "An improved bacterial colony lysis procedure enables direct DNA hybridisation using short (10, 11 bases) oligonucleotides to cosmids," *Nuc. Acids Res.*, 19(1):182 (1990).
- Nizetic et al., "Construction, arraying, and high-density screening of large insert libraries of human chromosomes X and 21: their potential use as reference libraries," *PNAS*, 88:3233-3237 (1991).
- Nyborg, W., "Acoustic Streaming," chapter 11 pp. 265-329 from *Physical Acoustics, Principles and Methods*, Mason, eds., vol. II, part B, Academic Press, New York and London (1965).
- Ocvirk et al., "High Performance Liquid Chromatography Partially Integrated onto a Silicon Chip," *Analyt. Meth. Instrumentation*, 2(2):74-82 (1995).
- Ohitsuka et al., "Studies on transfer ribonucleic acids and related compounds. IX Ribonucleic oligonucleotide synthesis using a photosensitive 0-nitrobenzyl protection at the 2'-hydroxyl group," *Nuc.Acids.Res.*, 1(10):1351-1357 (1974).
- Olefirowicz et al., "Capillary Electrophoresis for Sampling Single Nerve Cells," *Chimia*, 45(4):106-108 (1991).
- Olson et al., "Random-clone strategy for genomic restriction mapping in yeast," *PNAS*, 83:7826-7830 (1986).
- Patchornik et al., "Photosensitive Protecting Groups," *J.Am. Chem.Soc.*, 92(21):6333-6335 (1970).
- Patent Abstracts of Japan from EPO, Abst. 13:557, JP 1-233 447 (1989).
- Pease et al., "Light-generated oligonucleotide arrays for rapid DNA sequence analysis," *PNAS*, 91:5022-26 (1994).
- Pevzner, P.A., "DNA Physical Mapping and Alternating Eulerian Cycles in Colored Grapes," *Algorithmica*, 13(1-2):77-105 (1995).
- Pevzner et al., "Multiple Filtration and Approximate Pattern Matching," *Algorithmica*, 13(1-2):135-154 (1995).
- Pevzner et al., "Generalized Sequence Alignment and Duality," *Adv. Applied Math.*, 14:139-171 (1993).
- Pevzner, P.A., "1-Tuple DNA Sequencing: Computer Analysis," *J. Biomol. Struct. Dynam.*, 7(1):63-69 (1989).
- Pfahler et al., "Liquid Transport in Micron and Submicron Channels," *Sensors and Actuators*, A21-A23:431-4 (90).
- Pfeifer et al., "Genomic Sequencing and Methylation Analysis by Ligation Mediated PCR," *Science*, 246:810-813 (1989).
- Pidgeon et al., "Immobilized Artificial Membrane Chromatography: Supports Composed of Membrane Lipids," *Anal. Biochem.*, 176:36-47 (89).
- Pillai, V.N., "Photoremovable Protecting Groups in Organic Synthesis," *Synthesis*, pp. 1-26 (1980).
- Pillai et al., "3-Nitro-4-Aminomethylbenzoyl derivative von Polyethylenglykolen: Eine neue Klasse von Photosensitiven löslichen Polymeren Trägern zur Synthese von C-terminalen Peptidamiden," *Tetrah. Lett.*, # 36 p. 3409-3412 (1979).
- Pillai et al., "Synthesis Hydrophilic Polymers, Biomedical and Chemical Applications," *Naturwissenschaften*, 68:558-566 (1981).
- Pirrung et al., "Proofing of Photolithographic DNA Synthesis with 3'-5'-Dimethoxybenzoyloxycarbonyl-Protected Deoxynucleoside Phosphoramidites," *J. Org. Chem.*, 63(2):241-246 (1998).
- Pirrung et al., "Comparison of Methods for Photochemical Phosphoramidite-Based DNA Synthesis," *J. Org. Chem.*, 60:6270-6276 (1995).
- Ploax et al., "Cyclization of peptides on a solid support," *Int. J. Peptide Protein Research*, 29:162-169 (1987).
- Polsky-Cynkin et al., "Use of DNA Immobilized on Plastic and Agarose Supports to Detect DNA by Sandwich Hybridization," *Clin. Chem.*, 31(9):1428-1443 (1985).
- Poustka et al., "Molecular Approaches to Mammalian Genetics," Cold Spring Harbor Symposia on Quantitative Biology, 51:131-139 (1986).
- Purushothaman et al., "Synthesis of 4,5-diarylimidazole-2 thiones and their photoconversion to bis(4, 5-diarylimidazol-2-yl) sulphides," *Ind. J. Chem.*, 29B:18-21 (1990).
- Quesada et al., "High-Sensitivity DNA Detection with a Laser-Excited Confocal Fluorescence Gel Scanner," *Biotechniques*, 10:616 (1991).
- Reichmanis et al., *J. Polymer Sci. Polymer Chem. Edition*, 23:1-8 (1985).
- Renz et al., "A colorimetric method for DNA hybridization," *Nuc. Acids Res.*, 12(8):3435-3445 (1984).
- Richter et al., "An Electrohydrodynamic Micropump," *IEEE*, pp. 99-104 (1990).
- Richter et al., "Electrohydrodynamic Pumping and Flow Measurement," *IEEE*, pp. 271-276 (1991).
- Richter et al., "A Micromachined electrohydrodynamic (EHD) pump," *Sensors and Actuators*, A29:159-168 (91).
- Robertson et al., "A General and Efficient Route for Chemical Aminoacylation of Transfer RNAs," *J. Am. Chem. Soc.*, 113:2722-2729 (1991).
- Rodda et al., "The Antibody Response to Myoglobin-I. Systematic Synthesis of Myoglobin Peptides Reveals Location and Substructure of Species-Dependent Continuous Antigenic Determinants," *Mol. Immunol.*, 23(6):603-610 (1986).
- Rodgers, R.P., "Data Processing of Immunoassay Results," *Manual of Clin. Lab. Immunol.*, 3rd ed., ch. 15, pp. 82-87 (1986).
- Rose, D.J., "Free-solution reactor for post-column fluorescence detection in capillary zone electrophoresis," *J. Chromatography*, 540:343-353 (1991).
- Rovero et al., "Synthesis of Cyclic Peptides on solid Support," *Tetrahed. Letters*, 32(23):2639-2642 (1991).
- Sambrook, *Molecular Cloning—A Laboratory Manual*, publ. in 1989 (not included).
- Saiki et al., "Genetic analysis of amplified DNA with immobilized sequence-specific oligonucleotide probes," *PNAS*, 86:6230-6234 (1989).
- Saiki et al., "Analysis of enzymatically amplified  $\beta$ -globin and HLA-DQ $\alpha$  DNA with Allele-specific oligonucleotide probes," *Nature*, 324:163-166 (1986).
- Schafer et al., "DNA fingerprinting using non-radioactive oligonucleotide probes specific for simple repeats," *Nuc. Acids Res.*, 16(19):9344 (1988).
- Scharf et al., "HLA class II allelic variation and susceptibility to pemphigus vulgaris," *PNAS*, 85(10):3504-3508 (1988).

IAFP00000663

US 6,355,432 B1

Page 10

- Schena et al., "Parallel human genome analysis: Microarray-based expression monitoring of 1000 genes," *PNAS*, 93:10614-10619 (1996).
- Schuup et al., "Mechanistic Studies of the Photorearrangement of o-Nitrobenzyl Esters," *J. Photochem.*, 36:85-97 (1987).
- Seed, B., "Diazotizable arylamine cellulose papers for the coupling and hybridization of nucleic acids," *Nuc. Acids Res.*, 10(5):1799-1810 (1982).
- Seiler et al., "Planar Glass Chips for Capillary Electrophoresis: Repetitive Sample Injection, Quantitation, and Separation Efficiency," *Anal. Chem.*, 65:1481-1488 (1993).
- Seller et al., "Electroosmotic Pumping and Valveless Control of Fluid Flow with a Manifold of Capillaries on a Glass Chip," *Anal. Chem.*, 66:3485-3491 (1994).
- Semmelhack et al., "Selective Removal of Protecting Groups Using Controlled Potential Electrolysis," *J. Am. Chem. Society*, 94(14):5139-5140 (1972).
- Sheldon et al., "Matrix DNA Hybridization," *Clinical Chemistry*, 39(4):718-719 (1993).
- Shin et al., "Dehydrooligonopeptides. XI. Facile Synthesis of Various Kinds of Dipeptide- and tripeptides, and Dehydroenkephalins Containing Tyr Residue by Using N-Carboxydehydrotyrosine Anhydride," *Bull. Chem. Soc. Jpn.*, 62:1127-1135 (1989).
- Sim et al., "Use of a cDNA Library for Studies on Evolution and Developmental Expression of the Chorion Multigene Families," *Cell*, 18:1303-1316 (1979).
- Smith et al., "A Novel Method for Delineating Antigenic Determinants: Peptide Synthesis and Radioimmunoassay Using the Same Solid Support," *Immunochimistry*, 14:565-568 (1977).
- Sofia, M.J., "Carbohydrate-based combinatorial libraries," *Molecular Diversity*, 3:75-94 (1998).
- Southern et al., "Report on the Sequencing by Hybridization Workshop," *Genomics*, 13:1378-1383 (1992).
- Southern et al., "Oligonucleotide hybridisations on glass supports: a novel linker for oligonucleotide synthesis and hybridization properties of oligonucleotides synthesized in situ," *Nuc. Acids Res.*, 20(7):1679-1684 (1992).
- Southern et al., "Analyzing and Comparing Nucleic Acid Sequences by Hybridization to Arrays of Oligonucleotides: Evaluation Using Experimental Models," *Genomics*, 13:1008-10017 (1992).
- Southern, E.M., "Detection of Specific Sequences Among DNA Fragments Separated by Gel Electrophoresis," *J. Mol. Biol.*, 98:503-517 (1975).
- Stemme et al., "A valveless diffuser/nozzle-based fluid pump," *Sensors and Actuators*, A39:159-167 (1993).
- Stryer, L., "DNA Probes and Genes Can be Synthesized by Automated Solid-Phase Methods," from *Biochemistry*, Third Edition, published by W.H. Freeman & Co., (1988).
- Stuber et al., "Synthesis and photolytic cleavage of bovine insulin B22-30 on a nitrobenzoylglycyl-poly (ethylene glycol) support," *Int. J. Peptide Protein Res.*, 22(3):277-283 (1984).
- Sundberg et al., "Spatially-Addressable Immobilization of Macromolecules on Solid Supports," *J. Am. Chem. Soc.*, 117(49):12050-12057 (1995).
- Swedberg, S.A., "Use of non-ionic and zwitterionic surfactants to enhance selectivity in high-performance capillary electrophoresis, An apparent micellar electrokinetic capillary chromatography mechanism," *J. Chromatography*, 503:449-452 (1990).
- Thomas, P.S., "Hybridization of denatured RNA and small DNA fragments transferred to nitrocellulose," *PNAS*, 77(9):5201-5205 (1980).
- Titus et al., "Texas Red, a Hydrophilic, red-emitting fluorophore for use with fluorescein in dual parameter flow microfluorometric and fluorescence microscopic studies," *J. Immunol. Meth.*, 50:193-204 (1982).
- Tkachuk et al., "Detection of bcr-abl Fusion in chronic Myelogenous Leukemia by in situ Hybridization," *Science*, 250:559-562 (90).
- Trzeciak et al., "Synthesis of 'Head-to-Tail' Cyclized Peptides on Solid Support by Fmoc Chemistry," *Tetrahed. Letters*, 33(32):4557-4560 (1992).
- Tsien et al., "Control of Cytoplasmic Calcium with Photolabile Tetracarboxylate 2-Nitrobenzhydryl Chelators," *Biophys. J.*, 50:843-853 (1986).
- Tsutsumi et al., "Expression of L- and M- Type Pyruvate Kinase in Human Tissues," *Genomics*, 2:86-89 (1988).
- Turchinskii et al., "Multiple Hybridization in Genome Analysis, Reaction of Diamines and Bisulfate with Cytosine for Introduction of Nonradioactive Labels Into DNA," *Molecular Biology*, 22:1229-1235 (1988).
- Turner et al., "Photochemical Activation of Acylated  $\alpha$ -Thrombin," *J. Am. Chem. Soc.*, 109:1274-1275 (1987).
- Urdea et al., "A novel method for the rapid detection of specific nucleotide sequences in crude biological samples without blotting or radioactivity; application to the analysis of hepatitis B virus in human serum," *Gene*, 61:253-264 (1987).
- Urdea et al., "A comparison of non-radioisotopic hybridization assay methods using fluorescent, chemiluminescent and enzyme labeled synthetic oligodeoxyribonucleotide probes," *Nuc. Acids. Res.*, 16(11):4937-4956 (1988).
- Van der Voort et al., "Design and Use of a Computer Controlled Confocal Microscope for Biological Applications," *Scanning*, 7(2):66-78 (1985).
- Van Hijfte et al., "Intramolecular 1,3-Diyl Trapping Reactions. A Formal Total Synthesis of -Coriolin," *J. Organic Chemistry*, 50:3942-3944 (1985).
- Veldkamp, W.B., "Binary optics: the optics technology of the 1990s," *CLEO 90*, vol. 7, paper # CMG6 (1990).
- Verlaan-de Vries et al., "A dot-blot screening procedure for mutated ras oncogenes using synthetic oligodeoxynucleotides," *Gene*, 50:313-320 (1986).
- Verpoorte et al., "Three-dimensional micro flow manifolds for miniaturized chemical analysis systems," *J. Micromech. Microeng.*, 4:246-256 (1994).
- Volkmut et al., "DNA electrophoresis in microlithographic arrays," *Nature*, 358:600-602 (1992).
- Voss et al., "The immobilization of oligonucleotides and their hybridization properties," *Biochem. Soc. Transact.*, 16:216-217 (1988).
- Wada, A., *International Workshop on Automatic and High Speed DNA Base Sequencing*, Hayashibara Forum 1987 at Hayashibara Biochemical Laboratories, Okayama, Japan, Jul. 7-9, 1987.
- Walker et al., "Photolabile Protecting Groups for an Acetylcholine Receptor Ligand. Synthesis and Photochemistry of a New Class of o-Nitrobenzyl Derivatives and their Effects on Receptor Function," *Biochemistry*, 25:1799-1805 (1986).

IAFP00000664

US 6,355,432 B1

Page 11

- Wallace et al., "The use of synthetic oligonucleotides as hybridization probes. II. Hybridization of oligonucleotides of mixed sequence to rabbit  $\beta$ -globin DNA," *Nuc. Acids Res.*, 9(4):879 (1981).
- Wallace et al., "Hybridization of synthetic oligodeoxyribonucleotides to  $\Phi$ x 174 DNA: the effect of single base pair mismatch," *Nuc. Acids Res.*, 11(6):3543-3557 (1979).
- Washizu et al., "Handling Biological Cells Using a Fluid Integrated Circuit," *IEEE Transactions Industry Applications*, 26(2):352-358 (1990).
- Wiedmann, M. et al., "Ligase Chain Reaction (LCR)—Overview and Applications," *PCR Meth. Appl.*, 3(4):S51-S64 (1994).
- Werner et al., "Size-Dependent Separation of Proteins Denatured in SDS by Capillary Electrophoresis Using a Replaceable Sieving Matrix," *Anal. Biochem.*, 212:253-258 (1993).
- White et al., "An Evaluation of Confocal Versus Conventional Imaging of Biological Structures by Fluorescence Light Microscopy," *J. Cell Biol.*, 105(1):41-48 (1987).
- Widacki et al., "Biochemical Differences in Qa-2 Antigens Expressed by Qa-2+6+ and Qa-2a+6- Strains. Evidence for Differential Expression of the Q7 and Q9 Genes," *Mol. Immunology*, 27(6):559-570 (1990).
- Wilcox et al., "Synthesis of Photolabile 'Precursors' of Amino Acid Neurotransmitters," *J. Org. Chem.*, 55:1585-1589 (1990).
- Wilding et al., "PCR in a Silicon Microstructure," *Clin. Chem.*, 40(9):1815-1818 (1994).
- Wilding et al., "Manipulation and Flow of Biological Fluids in Straight Channels Micromachined in Silicon," *Clin. Chem.*, 40(1):43-47 (1994).
- Wittman-Liebold, eds., *Methods in Protein Sequence Analysis*, from Proceedings of 7th Int'l Conf., Berlin, Germany, 7/3-8/88, table of contents, pp. xi-xx\* (1989).
- Wood et al., "Base composition-independent hybridization in tetramethylammonium chloride: A method for oligonucleotide screening of highly complex gene libraries," *PNAS*, 82:1585-1588 (1985).
- Woolley et al., "Ultra-high-speed DNA fragment separations using microfabricated capillary array electrophoresis chips," *PNAS*, 91:11348-11352 (1994).
- Wu et al., "Synthesis and Properties of Adenosine-5'-triphospho- $\gamma$ -5-(5-sulfonic acid)naphthyl Ethylamide: A Fluorescent Nucleotide Substrate for DNA-Dependent RNA Polymerase from *Escherichia coli*," *Arch. Biochem. Biophys.*, 246(2):564-571 (1986).
- Wu et al., "Laboratory Methods, Direct Analysis of Single Nucleotide Variation in Human DNA and RNA Using In Situ Dot Hybridization," *DNA*, 8(2):135-142 (1989).
- Yamamoto et al., "Features and applications of the laser scanning microscope," *J. Mod. Optics*, 37(11):1691-1701 (1990).
- Yarbrough et al., "Synthesis and Properties of Fluorescent Nucleotide Substrates for DNA-dependent RNA Polymerases," *J. Biol. Chem.*, 254(23):12069-12073 (1979).
- Yosomiya et al., "Performance, Glass fiber Having Isocyanate Group on the Surface. Preparation and Reaction with Amino Acid," *Polymer Bulletin*, 12:41-48 (1984).
- Young, W.S., "Simultaneous Use of Digoxigenin- and Radiolabeled Oligodeoxyribonucleotide Probes for Hybridization Histochemistry," *Neuropeptides*, 13:271-275 (1989).
- Yue et al., "Miniature Field-Flow Fractionation System for Analysis of Blood Cells," *Clin. Chem.*, 40(9):1810-1814 (1994).
- Zehavi et al., "Light-Sensitive Glycosides. I. 6-Nitroveratryl  $\beta$ -D-Glucopyranoside and 2-Nitrobenzyl  $\beta$ -D-Glucopyranoside," *J. Org. Chem.*, 37(14):2281-2285 (1972).
- Zengerle et al., "Transient measurements on miniaturized diaphragm pumps in microfluid systems," *Sensors and Actuators*, A46-47:557-561 (1995).
- Zischler et al., "Non-radioactive oligonucleotide fingerprinting in the gel," *Nuc. Acids Res.*, 19(11):4411 (1989).
- Zischler et al., "Digoxigenated oligonucleotide probes specific for simple repeats in DNA fingerprinting and hybridization in situ," *Hum. Genet.*, 82:227-233 (1989).
- Hodgson et al., *Nucl. Acids Res.*, 15(15):6295 (1987).
- Khrapko et al., *DNA Seq. Map*, 1:375-388 (1991).
- Lander et al., *Genomics*, 2:231-239 (1988).
- Little, *Nature*, 346:611-612 (1990).
- Lysov et al., *Dokl. Akad. Nauk. SSSR*, 303:1508-1511 (1988).
- Olson et al., *Proc. Natl. Acad. Sci. USA*, 83:7826-7830 (Oct. 1986).
- Pevzner, *Algorithmica*, 13(1-2):77-105 (1995).
- Pevzner et al., *Algorithmica*, 13(1-2):135-154 (1995).
- Pfeifer et al., *Science*, 246:810-813 (Nov. 10, 1988).
- Seed, *Nucl. Acids Res.*, 10(5):1799-1810 (1982).
- Wood et al., *Proc. Natl. Acad. Sci. USA*, 82:1585-1588 (1985).
- Feinberg et al., *Anal. Biochem.*, 137:266-267 (1984).
- Pevzner et al., *Adv. Applied Math.*, 14:139-171 (1993).
- Schena et al., *Proc. Natl. Acad. Sci. USA*, 93:10614-10619 (Oct. 1996).
- Miller et al., "Detection of bacteria by hybridization of rRNA with DNA-latex and immunodetection of hybrids" *J Clin Microbiol* 1988, 26:1271-1276.
- Brenner et al., "In vitro cloning of complex mixtures of DNA on microbeads: Physical separation of differentially expressed cDNAs," *PNAS*, vol. 97, No. 4, Feb. 15, 2000, pp. 1665-1670.
- Brenner et al., "Gene expression analysis by massively parallel signature sequencing (MPSS) on microbead arrays," *Nature Biotechnology*, vol. 18, Jun. 2000, pp. 630-634.
- Tyagi, "Taking a census of mRNA populations with microbeads," *Nature Biotechnology*, vol. 18, Jun. 2000, pp. 597 and 598.

U.S. Patent

Mar. 12, 2002

Sheet 1 of 2

US 6,355,432 B1




FIG. 1



U.S. Patent

Mar. 12, 2002

Sheet 2 of 2

US 6,355,432 B1




FIG. 2A




FIG. 2B




FIG. 2C




FIG. 2D




FIG. 2E




FIG. 2F




FIG. 2G




FIG. 2H




FIG. 2I




FIG. 2K




FIG. 2L




FIG. 2M




FIG. 2N

US 6,355,432 B1

1

## PRODUCTS FOR DETECTING NUCLEIC ACIDS

### CROSS REFERENCE TO RELATED APPLICATIONS

This is a continuation of application Ser. No. 09/362,089, filed Jul. 28, 1998, pending; which is a divisional of application Ser. No. 09/056,927, filed Apr. 8, 1998, now U.S. Pat. No. 6,197,506; which is a continuation of application Ser. No. 08/670,118, filed Jun. 25, 1996, now U.S. Pat. No. 5,800,992; which is a divisional of application Ser. No. 08/168,904, filed Dec. 15, 1993, now abandoned; which is a continuation of application Ser. No. 07/624,114, filed Dec. 6, 1990, now abandoned; each of which is hereby incorporated by reference.

Additional commonly assigned application No. 07/492,462, filed Mar. 7, 1990, now U.S. Pat. No. 5,143,854; application No. 07/362,901, filed Jun. 7, 1989, now abandoned; application Ser. Nos. 07/624,120 and 07/626,730, both of which were filed on Dec. 6, 1990; application Ser. No. 07/435,316, filed Nov. 13, 1989, now abandoned; and U.S. Pat. No. 5,252,743 are also hereby incorporated herein by reference.

### BACKGROUND OF THE INVENTION

The present invention relates to the sequencing, fingerprinting, and mapping of polymers, particularly biological polymers. The inventions may be applied, for example, in the sequencing, fingerprinting, or mapping of nucleic acids, polypeptides, oligosaccharides, and synthetic polymers.

The relationship between structure and function of macromolecules is of fundamental importance in the understanding of biological systems. These relationships are important to understanding, for example, the functions of enzymes, structural proteins, and signalling proteins, ways in which cells communicate with each other, as well as mechanisms of cellular control and metabolic feedback.

Genetic information is critical in continuation of life processes. Life is substantially informationally based and its genetic content controls the growth and reproduction of the organism and its complements. Polypeptides, which are critical features of all living systems, are encoded by the genetic material of the cell. In particular, the properties of enzymes, functional proteins, and structural proteins are determined by the sequence of amino acids which make them up. As structure and function are integrally related, many biological functions may be explained by elucidating the underlying structural features which provide those functions. For this reason, it has become very important to determine the genetic sequences of nucleotides which encode the enzymes, structural proteins, and other effectors of biological functions. In addition to segments of nucleotides which encode polypeptides, there are many nucleotide sequences which are involved in control and regulation of gene expression.

The human genome project is directed toward determining the complete sequence of the genome of the human organism. Although such a sequence would not correspond to the sequence of any specific individual, it would provide significant information as to the general organization and specific sequences contained within segments from particular individuals. It would also provide mapping information which is very useful for further detailed studies. However, the need for highly rapid, accurate, and inexpensive sequencing technology is nowhere more apparent than in a

2

demanding sequencing project such as this. To complete the sequencing of a human genome would require the determination of approximately  $3 \times 10^9$ , or 3 billion base pairs.

The procedures typically used today for sequencing include the Sanger dideoxy method, see, e.g., Sanger et al. (1977) *Proc. Natl. Acad. Sci. USA*, 74:5463-5467, or the Maxam and Gilbert method, see, e.g., Maxam et al., (1980) *Methods in Enzymology*, 65:499-559. The Sanger method utilizes enzymatic elongation procedures with chain terminating nucleotides. The Maxam and Gilbert method uses chemical reactions exhibiting specificity of reaction to generate nucleotide specific cleavages. Both methods require a practitioner to perform a large number of complex manual manipulations. These manipulations usually require isolating homogeneous DNA fragments, elaborate and tedious preparing of samples, preparing a separating gel, applying samples to the gel, electrophoresing the samples into this gel, working up the finished gel, and analyzing the results of the procedure.

Thus, a less expensive, highly reliable, and labor efficient means for sequencing biological macromolecules is needed. A substantial reduction in cost and increase in speed of nucleotide sequencing would be very much welcomed. In particular, an automated system would improve the reproducibility and accuracy of procedures. The present invention satisfies these and other needs.

### SUMMARY OF THE INVENTION

The present invention provides improved methods useful for de novo sequencing of an unknown polymer sequence, for verification of known sequences, for fingerprinting polymers, and for mapping homologous segments within a sequence. By reducing the number of manual manipulations required and automating most of the steps, the speed, accuracy, and reliability of these procedures are greatly enhanced.

The production of a substrate having a matrix of positionally defined regions with attached reagents exhibiting known recognition specificity can be used for the sequence analysis of a polymer. Although most directly applicable to sequencing, the present invention is also applicable to fingerprinting, mapping, and general screening of specific interactions. The VLSIPS™ Technology (Very Large Scale Immobilized Polymer Synthesis) substrates will be applied to evaluating other polymers, e.g., carbohydrates, polypeptides, hydrocarbon synthetic polymers, and the like. For these non-polynucleotides, the sequence specific reagents will usually be antibodies specific for a particular subunit sequence.

According to one aspect of the masking technique, the invention provides an ordered method for forming a plurality of polymer sequences by sequential addition of reagents comprising the step of serially protecting and deprotecting portions of the plurality of polymer sequences for addition of other portions of the polymer sequences using a binary synthesis strategy.

The present invention also provides a means to automate sequencing manipulations. The automation of the substrate production method and of the scan and analysis steps minimizes the need for human intervention. This simplifies the tasks and promotes reproducibility.

The present invention provides a composition comprising a plurality of positionally distinguishable sequence specific reagents attached to a solid substrate, which reagents are capable of specifically binding to a predetermined subunit sequence of a preselected multi-subunit length having at

US 6,355,432 B1

3

least three subunits, said reagents representing substantially all possible sequences of said preselected length. In some embodiments, the subunit sequence is a polynucleotide or a polypeptide, in others the preselected multi-subunit length is five subunits and the subunit sequence is a polynucleotide sequence. In other embodiments, the specific reagent is an oligonucleotide of at least about five nucleotides. Alternatively, the specific reagent is a monoclonal antibody. Usually the specific reagents are all attached to a single solid substrate, and the reagents comprise about 3000 different sequences. In other embodiments, the reagents represents at least about 25% of the possible subsequences of said preselected length. Usually, the reagents are localized in regions of the substrate having a density of at least 25 regions per square centimeter, and often the substrate has a surface area of less than about 4 square centimeters.

The present invention also provides methods for analyzing a sequence of a polynucleotide or a polypeptide, said method comprising the step of:

- a) exposing said polynucleotide or polypeptide to a composition as described.

It also provides useful methods for identifying or comparing a target sequence with a reference, said method comprising the step of:

- a) exposing said target sequence to a composition as described;
- b) determining the pattern of positions of the reagents which specifically interact with the target sequence; and
- c) comparing the pattern with the pattern exhibited by the reference when exposed to the composition.

The present invention also provides methods for sequencing a segment of a polynucleotide comprising the steps of:

- a) combining:
  - i) a substrate comprising a plurality of chemically synthesized and positionally distinguishable oligonucleotides capable of recognizing defined oligonucleotide sequences; and
  - ii) a target polynucleotide; thereby forming high fidelity matched duplex structures of complementary subsequences of known sequence; and
- b) determining which of said reagents have specifically interacted with subsequences in said target polynucleotide.

In one embodiment, the segment is substantially the entire length of said polynucleotide.

The invention also provides methods for sequencing a polymer, said method comprising the steps of:

- a) preparing a plurality of reagents which each specifically bind to a subsequence of preselected length;
- b) positionally attaching each of said reagents to one or more solid phase substrates, thereby producing substrates of positionally definable sequence specific probes;
- c) combining said substrates with a target polymer whose sequence is to be determined; and
- d) determining which of said reagents have specifically interacted with subsequences in said target polymer.

In one embodiment, the substrates are beads. Preferably, the plurality of reagents comprise substantially all possible subsequences of said preselected length found in said target. In another embodiment, the solid phase substrate is a single substrate having attached thereto reagents recognizing substantially all possible subsequences of preselected length found in said target.

4

In another embodiment, the method further comprises the step of analyzing a plurality of said recognized subsequences to assemble a sequence of said target polymer. In a bead embodiment, at least some of the plurality of substrates have one subsequence specific reagent attached thereto, and the substrates are coded to indicate the sequence specificity of said reagent.

The present invention also embraces a method of using a fluorescent nucleotide to detect interactions with oligonucleotide probes of known sequence, said method comprising:

- a) attaching said nucleotide to a target unknown polynucleotide sequence, and
- b) exposing said target polynucleotide sequence to a collection of positionally defined oligonucleotide probes of known sequences to determine the sequences of said probes which interact with said target.

In a further refinement, an additional step is included of:

- a) collating said known sequences to determine the overlaps of said known sequences to determine the sequence of said target sequence.

A method of mapping a plurality of sequences relative to one another is also provided, the method comprising:

- a) preparing a substrate having a plurality of positionally attached sequence specific probes;
- b) exposing each of said sequences to said substrate, thereby determining the patterns of interaction between said sequence specific probes and said sequences; and
- c) determining the relative locations of said sequence specific probe interactions on said sequences to determine the overlaps and order of said sequences.

In one refinement, the sequence specific probes are oligonucleotides, applicable to where the target sequences are nucleic acid sequences.

In the nucleic acid sequencing application, the steps of the sequencing process comprise:

- a) producing a matrix substrate having known positionally defined regions of known sequence specific oligonucleotide probes;
- b) hybridizing a target polynucleotide to the positions on the matrix so that each of the positions which contain oligonucleotide probes complementary to a sequence on the target hybridize to the target molecule;
- c) detecting which positions have bound the target, thereby determining sequences which are found on the target; and
- d) analyzing the known sequences contained in the target to determine sequence overlaps and assembling the sequence of the target therefrom.

The enablement of the sequencing process by hybridization is based in large part upon the ability to synthesize a large number (e.g., to virtually saturate) of the possible overlapping sequence segments and distinguishing those probes which hybridize with fidelity from those which have mismatched bases, and to analyze a highly complex pattern of hybridization results to determine the overlap regions.

The detecting of the positions which bind the target sequence would typically be through a fluorescent label on the target. Although a fluorescent label is probably most convenient, other sorts of labels, e.g., radioactive, enzyme linked, optically detectable, or spectroscopic labels may be used. Because the oligonucleotide probes are positionally defined, the location of the hybridized duplex will directly translate to the sequences which hybridize. Thus, analysis of the positions provides a collection of subsequences found within the target sequence. These subsequences are matched

US 6,355,432 B1

5

with respect to their overlaps so as to assemble an intact target sequence.

#### BRIEF DESCRIPTION OF THE DRAWINGS

FIG. 1 illustrates a flow chart for sequence, fingerprint, or mapping analysis.

FIGS. 2A-2M illustrate the proper function of a VLSIPS™ Technology nucleotide synthesis.

#### DESCRIPTION OF THE PREFERRED EMBODIMENTS

- I. Overall Description
  - A. general
  - B. VLSIPS substrates
  - C. binary masking
  - D. applications
  - E. detection methods and apparatus
  - F. data analysis
- II. Theoretical Analysis
  - A. simple n-mer structure; theory
  - B. complications
  - C. non-polynucleotide embodiments
- III. Polynucleotide Sequencing
  - A. preparation of substrate matrix
  - B. labeling target polynucleotide
  - C. hybridization conditions
  - D. detection; VLSIPS scanning
  - E. analysis
  - F. substrate reuse
  - G. non-polynucleotide aspects
- IV. Fingerprinting
  - A. general
  - B. preparation of substrate matrix
  - C. labeling target nucleotides
  - D. hybridization conditions
  - E. detection; VLSIPS scanning
  - F. analysis
  - G. substrate reuse
  - H. non-polynucleotide aspects
- V. Mapping
  - A. general
  - B. preparation of substrate matrix
  - C. labeling
  - D. hybridization/specific interaction
  - E. detection
  - F. analysis
  - G. substrate reuse
  - H. non-polynucleotide aspects
- VI. Additional Screening
  - A. specific interactions
  - B. sequence comparisons
  - C. categorizations
  - D. statistical correlations
- VII. Formation of Substrate
  - A. instrumentation
  - B. binary masking
  - C. synthetic methods
  - D. surface immobilization

6

#### VIII. Hybridization/Specific Interaction

- A. general
- B. important parameters

#### IX. Detection Methods

- A. labeling techniques
- B. scanning system

#### X. Data Analysis

- A. general
- B. hardware
- C. software

#### XI. Substrate Reuse

- A. removal of label
- B. storage and preservation

#### XII. Integrated Sequencing Strategy

- A. initial mapping strategy
- B. selection of smaller clones
- C. actual sequencing procedures

#### XIII. Commercial Applications

- A. sequencing
- B. fingerprinting
- C. mapping

#### I. OVERALL DESCRIPTION

##### A. General

The present invention relies in part on the ability to synthesize or attach specific recognition reagents at known locations on a substrate, typically a single substrate. In particular, the present invention provides the ability to prepare a substrate having a very high density matrix pattern of positionally defined specific recognition reagents. The reagents are capable of interacting with their specific targets while attached to the substrate, e.g., solid phase interactions, and by appropriate labeling of these targets, the sites of the interactions between the target and the specific reagents may be derived. Because the reagents are positionally defined, the sites of the interactions will define the specificity of each interaction. As a result, a map of the patterns of interactions with specific reagents on the substrate is convertible into information on the specific interactions taking place, e.g., the recognized features. Where the specific reagents recognize a large number of possible features, this system allows the determination of the combination of specific interactions which exist on the target molecule. Where the number of features is sufficiently large, the identical same combination, or pattern, of features is sufficiently unlikely that a particular target molecule may often be uniquely defined by its features. In the extreme, the features may actually be the subunit sequence of the target molecule, and a given target sequence may be uniquely defined by its combination of features.

In particular, the methodology is applicable to sequencing polynucleotides. The specific sequence recognition reagents will typically be oligonucleotide probes which hybridize with specificity to subsequences found on the target sequence. A sufficiently large number of those probes allows the fingerprinting of a target polynucleotide or the relative mapping of a collection of target polynucleotides, as described in greater detail below.

In the high resolution fingerprinting provided by a saturating collection of probes which include all possible subsequences of a given size, e.g., 10-mers, collating of all the subsequences and determination of specific overlaps will be derived and the entire sequence can usually be reconstructed.



US 6,355,432 B1

7

Although a polynucleotide sequence analysis is a preferred embodiment, for which the specific reagents are most easily accessible, the invention is also applicable to analysis of other polymers, including polypeptides, carbohydrates, and synthetic polymers, including  $\alpha$ -,  $\beta$ -, and  $\omega$ -amino acids, polyurethanes, polyesters, polycarbonates, polyureas, polyamides, polyethyleneimines, polyarylene sulfides, polysiloxanes, polyimides, polyacetates, and mixed polymers. Various optical isomers, e.g., various D- and L-forms of the monomers, may be used.

Sequence analysis will take the form of complete sequence determination, to the level of the sequence of individual subunits along the entire length of the target sequence. Sequence analysis also takes the form of sequence homology, e.g., less than absolute subunit resolution, where "similarity" in the sequence will be detectable, or the form of selective sequences of homology interspersed at specific or irregular locations.

In either case, the sequence is determinable at selective resolution or at particular locations. Thus, the hybridization method will be useful as a means for identification, e.g., a "fingerprint", much like a Southern hybridization method is used. It is also useful to map particular target sequences.

#### B. VLSIPS™ Technology

The invention is enabled by the development of technology to prepare substrates on which specific reagents may be either positionally attached or synthesized. In particular, the very large scale immobilized polymer synthesis (VLSIPS™) technology allows for the very high density production of an enormous diversity of reagents mapped out in a known matrix pattern on a substrate. These reagents specifically recognize subsequences in a target polymer and bind thereto, producing a map of positionally defined regions of interaction. These map positions are convertible into actual features recognized, and thus would be present in the target molecule of interest.

As indicated, the sequence specific recognition reagents will often be oligonucleotides which hybridize with fidelity and discrimination to the target sequence. For use with other polymers, monoclonal or polyclonal antibodies having high sequence specificity will often be used.

In the genetic sense, the VLSIPS technology allows the production of a substrate with a high density matrix of positionally mapped regions with specific recognition reagents attached at each distinct region. By use of protective groups which can be positionally removed, or added, the regions can be activated or deactivated for addition of particular reagents or compounds. Details of the protection are described below and in related Pirrung et al. (1992) U.S. Pat. No. 5,143,854. In a preferred embodiment, photosensitive protecting agents will be used and the regions of activation or deactivation may be controlled by electro-optical and optical methods, similar to many of the processes used in semiconductor wafer and chip fabrication.

In the nucleic acid nucleotide sequencing application, a VLSIPS substrate is synthesized having positionally defined oligonucleotide probes. See Pirrung et al. (1992) U.S. Pat. No. 5,143,854; and U.S. Ser. No. 07/624,120, now abandoned. By use of masking technology and photosensitive synthetic subunits, the VLSIPS apparatus allows for the stepwise synthesis of polymers according to a positionally defined matrix pattern. Each oligonucleotide probe will be synthesized at known and defined positional locations on the substrate. This forms a matrix pattern of known relationship between position and specificity of interaction. The VLSIPS technology allows the production of a very large number of different oligonucleotide probes to be simultaneously and

8

automatically synthesized including numbers in excess of about  $10^2$ ,  $10^3$ ,  $10^4$ ,  $10^5$ ,  $10^6$ , or even more, and at densities of at least about  $10^2$ ,  $10^3/\text{cm}^2$ ,  $10^4/\text{cm}^2$ ,  $10^5/\text{cm}^2$  and up to  $10^6/\text{cm}^2$  or more. This application discloses methods for synthesizing polymers on a silicon or other suitably derivatized substrate, methods and chemistry for synthesizing specific types of biological polymers on those substrates, apparatus for scanning and detecting whether interaction has occurred at specific locations on the substrate, and various other technologies related to the use of a high density very large scale immobilized polymer substrate. In particular, sequencing, fingerprinting, and mapping applications are discussed herein in detail, though related technologies are described in simultaneously filed applications U.S. Ser. No. 07/624,120, now abandoned; and U.S. Ser. No. 07/517,659; Dower et al. (1995) U.S. Pat. No. 5,427,908, each of which is hereby incorporated herein by reference.

In other embodiments, antibody probes will be generated which specifically recognize particular subsequences found on a polymer. Antibodies would be generated which are specific for recognizing a three contiguous amino acid sequence, and monoclonal antibodies may be preferred. optimally, these antibodies would not recognize any sequences other than the specific three amino acid stretch desired and the binding affinity should be insensitive to flanking or remote sequences found on a target molecule. Likewise, antibodies specific for particular carbohydrate linkages or sequences will be generated. A similar approach could be used for preparing specific reagents which recognize other polymer subunit sequences. These reagents would typically be site specifically localized to a substrate matrix pattern where the regions are closely packed.

These reagents could be individually attached at specific sites on the substrate in a matrix by an automated procedure where the regions are positionally targeted by some other specific mechanism, e.g., one which would allow the entire collection of reagents to be attached to the substrate in a single reaction. Each reagent could be separately attached to a specific oligonucleotide sequence by an automated procedure. This would produce a collection of reagents where, e.g., each monoclonal antibody would have a unique oligonucleotide sequence attached to it. By virtue of a VLSIPS substrate which has different complementary oligonucleotides synthesized on it, each monoclonal antibody would specifically be bound only at that site on the substrate where the complementary oligonucleotide has been synthesized. A crosslinking step would fix the reagent to the substrate. See, e.g., Dattagupta et al. (1985) U.S. Pat. No. 4,542,102 and (1987) U.S. Pat. No. 4,713,326; and Chatterjee, M. et al. (1990) *J. Am. Chem. Soc.* 112:6397-6399, which are hereby incorporated herein by reference. This allows a high density positionally specific collection of specific recognition reagents, e.g., monoclonal antibodies, to be immobilized to a solid substrate using an automated system.

The regions which define particular reagents will usually be generated by selective protecting groups which may be activated or deactivated. Typically the protecting group will be bound to a monomer subunit or spatial region, and can be spatially affected by an activator, such as electromagnetic radiation. Examples of protective groups with utility herein include nitroveratryl oxycarbonyl (NVOC), nitrobenzyl oxycarbonyl (NBOC), dimethyl dimethoxy benzyl oxycarbonyl, 5-bromo-7-nitroindolyl, O-hydroxy- $\alpha$ -methyl cinnamoyl, and 2-oxymethylene anthraquinone. Examples of activators include ion beams, electric fields, magnetic fields, electron beams, x-ray, and other forms of electromagnetic radiation.



US 6,355,432 B1

9

## C. Binary Masking

In fact, the means for producing a substrate useful for these techniques are explained in Pirrung et al. (1992) U.S. Pat. No. 5,143,854, which is hereby incorporated herein by reference. However, there are various particular ways to optimize the synthetic processes. Many of these methods are described in Ser. No. 07/624,120, now abandoned.

Briefly, the binary synthesis strategy refers to an ordered strategy for parallel synthesis of diverse polymer sequences by sequential addition of reagents which may be represented by a reactant matrix, and a switch matrix, the product of which is a product matrix. A reactant matrix is a  $1 \times n$  matrix of the building blocks to be added. The switch matrix is all or a subset of the binary numbers from 1 to  $n$  arranged in columns. In preferred embodiments, a binary strategy is one in which at least two successive steps illuminate half of a region of interest on the substrate. In most preferred embodiments, binary synthesis refers to a synthesis strategy which also factors a previous addition step. For example, a strategy in which a switch matrix for a masking strategy halves regions that were previously illuminated, illuminating about half of the previously illuminated region and protecting the remaining half (while also protecting about half of previously protected regions and illuminating about half of previously protected regions). It will be recognized that binary rounds may be interspersed with non-binary rounds and that only a portion of a substrate may be subjected to a binary scheme, but will still be considered to be a binary masking scheme within the definition herein. A binary "masking" strategy is a binary synthesis which uses light to remove protective groups from materials for addition of other materials such as nucleotides or amino acids.

In particular, this procedure provides a simplified and highly efficient method for saturating all possible sequences of a defined length polymer. This masking strategy is also particularly useful in producing all possible oligonucleotide sequence probes of a given length.

## D. Applications

The technology provided by the present invention has very broad applications. Although described specifically for polynucleotide sequences, similar sequencing, fingerprinting, mapping, and screening procedures can be applied to polypeptide, carbohydrate, or other polymers. In particular, the present invention may be used to completely sequence a given target sequence to subunit resolution. This may be for de novo sequencing, or may be used in conjunction with a second sequencing procedure to provide independent verification. See, e.g., (1988) *Science* 242:1245. For example, a large polynucleotide sequence defined by either the Maxam and Gilbert technique or by the Sanger technique may be verified by using the present invention.

In addition, by selection of appropriate probes, a polynucleotide sequence can be fingerprinted. Fingerprinting is a less detailed sequence analysis which usually involves the characterization of a sequence by a combination of defined features. Sequence fingerprinting is particularly useful because the repertoire of possible features which can be tested is virtually infinite. Moreover, the stringency of matching is also variable depending upon the application. A Southern Blot analysis may be characterized as a means of simple fingerprint analysis.

Fingerprinting analysis may be performed to the resolution of specific nucleotides, or may be used to determine homologies, most commonly for large segments. In particular, an array of oligonucleotide probes of virtually any workable size may be positionally localized on a matrix and used to probe a sequence for either absolute comple-

10

mentary matching, or homology to the desired level of stringency using selected hybridization conditions.

In addition, the present invention provides means for mapping analysis of a target sequence or sequences. Mapping will usually involve the sequential ordering of a plurality of various sequences, or may involve the localization of a particular sequence within a plurality of sequences. This may be achieved by immobilizing particular large segments onto the matrix and probing with a shorter sequence to determine which of the large sequences contain that smaller sequence. Alternatively, relatively shorter probes of known or random sequence may be immobilized to the matrix and a map of various different target sequences may be determined from overlaps. Principles of such an approach are described in some detail by Evans et al. (1989) "Physical Mapping of Complex Genomes by Cosmid Multiplex Analysis," *Proc. Natl. Acad. Sci. USA* 86:5030-5034; Michiels et al. (1987) "Molecular Approaches to Genome Analysis: A Strategy for the Construction of Ordered Overlap Clone Libraries," *CABIOS* 3:203-210; Olsen et al. (1986) "Random-Clone Strategy for Genomic Restriction Mapping in Yeast," *Proc. Natl. Acad. Sci. USA* 83:7826-7830; Craig, et al. (1990) "Ordering of Cosmid Clones Covering the Herpes Simplex Virus Type I (HSV-1) Genome: A Test Case for Fingerprinting by Hybridization," *Nuc. Acids Res.* 18:2653-2660; and Coulson, et al. (1986) "Toward a Physical Map of the Genome of the Nematode *Caenorhabditis elegans*," *Proc. Natl. Acad. Sci. USA* 83:7821-7825; each of which is hereby incorporated herein by reference.

Fingerprinting analysis also provides a means of identification. In addition to its value in apprehension of criminals from whom a biological sample, e.g., blood, has been collected, fingerprinting can ensure personal identification for other reasons. For example, it may be useful for identification of bodies in tragedies such as fire, flood, and vehicle crashes. In other cases the identification may be useful in identification of persons suffering from amnesia, or of missing persons. Other forensics applications include establishing the identity of a person, e.g., military identification "dog tags", or may be used in identifying the source of particular biological samples. Fingerprinting technology is described, e.g., in Carrano, et al. (1989) "A High-Resolution, Fluorescence-Based, Semi-automated method for DNA Fingerprinting," *Genomics* 4: 129-136, which is hereby incorporated herein by reference. See, e.g., table I, for nucleic acid applications, and corresponding applications may be accomplished using polypeptides.

TABLE I

## VLSIPS™ TECHNOLOGY IN NUCLEIC ACIDS

- |     |  |
|-----|--|
| I.  | Construction of Chips  |
| II. | Applications   |
| A.  | Sequencing   |
| 1.  | Primary sequencing   |
| 2.  | Secondary sequencing (sequence checking)                         |
| 3.  | Large scale mapping  |
| 4.  | Fingerprinting   |
| B.  | Duplex/Triplex formation   |
| 1.  | Antisense  |
| 2.  | Sequence specific function modulation (e.g. promoter inhibition) |
| C.  | Diagnosis  |
| 1.  | Genetic markers  |
| 2.  | Type markers   |
| a.  | Blood donors   |
| b.  | Tissue transplants   |

US 6,355,432 B1

11

TABLE I-continued

| VLSIPS™ TECHNOLOGY IN NUCLEIC ACIDS |  |
|-------------------------------------|--|
| D. Microbiology                     |  |
| 1. Clinical microbiology            |  |
| 2. Food microbiology                |  |
| III. Instrumentation                |  |
| A. Chip machines                    |  |
| B. Detection                        |  |
| IV. Software Development            |  |
| A. Instrumentation software         |  |
| B. Data reduction software          |  |
| C. Sequence analysis software       |  |

The fingerprinting analysis may be used to perform various types of genetic screening. For example, a single substrate may be generated with a plurality of screening probes, allowing for the simultaneous genetic screening for a large number of genetic markers. Thus, prenatal or diagnostic screening can be simplified, economized, and made more generally accessible.

In addition to the sequencing, fingerprinting, and mapping applications, the present invention also provides means for determining specificity of interaction with particular sequences. Many of these applications were described in Ser. No. 07/362,901, now abandoned, Pirrung et al. (1992) U.S. Pat. No. 5,143,854; Ser. No. 07/435,316, and Ser. No. 07/612,671.

#### E. Detection Methods and Apparatus

An appropriate detection method applicable to the selected labeling method can be selected. Suitable labels include radionucleotides, enzymes, substrates, cofactors, inhibitors, magnetic particles, heavy metal atoms, and particularly fluorescers, chemiluminescers, and spectroscopic labels. Patents teaching the use of such labels include U.S. Pat. Nos. 3,817,837; 3,850,752; 3,939,350; 3,996,345; 4,277,437; 4,275,149; and 4,366,241.

With an appropriate label selected, the detection system best adapted for high resolution and high sensitivity detection may be selected. As indicated above, an optically detectable system, e.g., fluorescence or chemiluminescence would be preferred. Other detection systems may be adapted to the purpose, e.g., electron microscopy, scanning electron microscopy (SEM), scanning tunneling electron microscopy (STEM), infrared microscopy, atomic force microscopy (AFM), electrical conductance, and image plate transfer.

With a detection method selected, an apparatus for scanning the substrate will be designed. Apparatus, as described in Ser. No. 07/362,901, now abandoned; or Pirrung et al. (1992) U.S. Pat. No. 5,143,854; or Ser. No. 07/624,120, now abandoned, are particularly appropriate. Design modifications may also be incorporated therein.

#### F. Data Analysis

Data is analyzed by processes similar to those described below in the section describing theoretical analysis. More efficient algorithms will be mathematically devised, and will usually be designed to be performed on a computer. Various computer programs which may more quickly or efficiently make measurement samples and distinguish signal from noise will also be devised. See, particularly, Ser. No. 07/624,120, now abandoned.

The initial data resulting from the detection system is an array of data indicative of fluorescent intensity versus location on the substrate. The data are typically taken over regions substantially smaller than the area in which synthesis of a given polymer has taken place. Merely by way of example, if polymers were synthesized in squares on the substrate having dimensions of 500 microns by 500 microns, the data may be taken over regions having dimensions of 5 microns by 5 microns. In most preferred embodiments, the

12

regions over which fluorescence data are taken across the substrate are less than about  $\frac{1}{2}$  the area of the regions in which individual polymers are synthesized, preferably less than  $\frac{1}{10}$  the area in which a single polymer is synthesized, and most preferably less than  $\frac{1}{100}$  the area in which a single polymer is synthesized. Hence, within any area in which a given polymer has been synthesized, a large number of fluorescence data points are collected.

A plot of number of pixels versus intensity for a scan should bear a rough resemblance to a bell curve, but spurious data are observed, particularly at higher intensities. Since it is desirable to use an average of fluorescent intensity over a given synthesis region in determining relative binding affinity, these spurious data will tend to undesirably skew the data.

Accordingly, in one embodiment of the invention the data are corrected for removal of these spurious data points, and an average of the data points is thereafter utilized in determining relative binding efficiency. In general the data are fitted to a base curve and statistical measures are used to remove spurious data.

In an additional analytical tool, various degeneracy reducing analogues may be incorporated in the hybridization probes. Various aspects of this strategy are described, e.g., in Macevitz, S. (1990) PCT publication number WO 90/04652, which is hereby incorporated herein by reference.

## II. THEORETICAL ANALYSIS

The principle of the hybridization sequencing procedure is based, in part, upon the ability to determine overlaps of short segments. The VLSIPS technology provides the ability to generate reagents which will saturate the possible short subsequence recognition possibilities. The principle is most easily illustrated by using a binary sequence, such as a sequence of zeros and ones. Once having illustrated the application to a binary alphabet, the principle may easily be understood to encompass three letter, four letter, five or more letter, even 20 letter alphabets. A theoretical treatment of analysis of subsequence information to reconstruction of a target sequence is provided, e.g., in Lysov, Yu., et al. (1988) *Doklady Akademi. Nauk. SSR* 303:1508-1511; Khrapko K., et al. (1989) *FEBS Letters* 256:118-122; Pevzner, P. (1989) *J. of Biomolecular Structure and Dynamics* 7:63-69; and Drmanac, R. et al. (1989) *Genomics* 4:114-128; each of which is hereby incorporated herein by reference.

The reagents for recognizing the subsequences will usually be specific for recognizing a particular polymer subsequence anywhere within a target polymer. It is preferable that conditions may be devised which allow absolute discrimination between high fidelity matching and very low levels of mismatching. The reagent interaction will preferably exhibit no sensitivity to flanking sequences, to the subsequence position within the target, or to any other remote structure within the sequence. For polynucleotide sequencing, the specific reagents can be oligonucleotide probes; for polypeptides and carbohydrates, antibodies will be useful reagents. Antibody reagents should also be useful for other types of polymers.

#### A. Simple n-mer Structure: Theory

##### 1. Simple Two Letter Alphabet: Example

A simple example is presented below of how a sequence of ten digits comprising zeros and ones would be sequenceable using short segments of five digits. For example, consider the sample ten digit sequence:

1010011100.

A VLSIPS™ Technology substrate could be constructed, as discussed elsewhere, which would have reagents attached in

US 6,355,432 B1

13

a defined matrix pattern which specifically recognize each of the possible five digit sequences of ones and zeros. The number of possible five digit subsequences is  $2^5=32$ . The number of possible different sequences 10 digits long is  $2^{10}=1,024$ . The five contiguous digit subsequences within a ten digit sequence number six, i.e., positioned at digits 1-5, 2-6, 3-7, 4-8, 5-9, and 6-10. It will be noted that the specific order of the digits in the sequence is important and that the order is directional, e.g., running left to right versus right to left. The first five digit sequence contained in the target sequence is 10100. The second is 01001, the third is 10011, the fourth is 00111, the fifth is 01110, and the sixth is 11100.

The VLSIPS™ substrate would have a matrix pattern of positionally attached reagents which recognize each of the different 5-mer subsequences. Those reagents which recognize each of the 6 contained 5-mers will bind the target, and a label allows the positional determination of where the sequence specific interaction has occurred. By correlation of the position in the matrix pattern, the corresponding bound subsequences can be determined.

In the above-mentioned sequence, six different 5-mer sequences would be determined to be present. They would be:

```

10100
01001
10011
00111
01110
11100

```

Any sequence which contains the first five digit sequence, 10100, already narrows the number of possible sequences (e.g., from 1024 possible sequences) which contain it to less than about 192 possible sequences.

This 192 is derived from the observation that with the subsequence 10100 at the far left of the sequence, in positions 1-5, there are only 32 possible sequences. Likewise, for that particular subsequence in positions 2-6, 3-7, 4-8, 5-9, and 6-10. So, to sum up all of the sequences that could contain 10100, there are 32 for each position and 6 positions for a total of about 192 possible sequences. However, some of these 10 digit sequences will have been counted twice. Thus, by virtue of containing the 10100 subsequence, the number of possible 10-mer sequences has been decreased from 1024 sequences to less than about 192 sequences.

In this example, not only do we know that the sequence contains 10100, but we also know that it contains the second five character sequence, 01001. By virtue of knowing that the sequence contains 10100, we can look specifically to determine whether the sequence contains a subsequence of five characters which contains the four leftmost digits plus a next digit to the left. For example, we would look for a sequence of X1010, but we find that there is none. Thus, we know that the 10100 must be at the left end of the 10-mer. We would also look to see whether the sequence contains the rightmost four digits plus a next digit to the right, e.g., 0100X. We find that the sequence also contains the sequence 01001, and that X is a 1. Thus, we know at least that our target sequence has an overlap of 0100 and has the left terminal sequence 101001.

Applying the same procedure to the second 5-mer, we also know that the sequence must include a sequence of five

14

digits having the sequence 1001Y where Y must be either 0 or 1. We look through the fragments and we see that we have a 10011 sequence within our target, thus Y is also 1. Thus, we would know that our sequence has a sequence of the first seven being 1010011.

Moving to the next 5-mer, we know that there must be a sequence of 0011Z, where Z must be either 0 or 1. We look at the fragments produced above and see that the target sequence contains a 00111 subsequence and Z is 1. Thus, we know the sequence must start with 10100111.

The next 5-mer must be of the sequence 0111W where W must be 0 or 1. Again, looking up at the fragments produced, we see that the target sequence contains a 01110 subsequence, and W is a 0. Thus, our sequence to this point is 101001110. We know that the last 5-mer must be either 11100 or 11101. Looking above, we see that it is 11100 and that must be the last of our sequence. Thus, we have determined that our sequence must have been 1010011100.

However, it will be recognized from the example above with the sequences provided therein, that the sequence analysis can start with any known positive probe subsequence. The determination may be performed by moving linearly along the sequence checking the known sequence with a limited number of next positions. Given this possibility, the sequence may be determined, besides by scanning all possible oligonucleotide probe positions, by specifically looking only where the next possible positions would be. This may increase the complexity of the scanning but may provide a longer time span dedicated towards scanning and detecting specific positions of interest relative to other sequence possibilities. Thus, the scanning apparatus could be set up to work its way along a sequence from a given contained oligonucleotide to only look at those positions on the substrate which are expected to have a positive signal.

It is seen that given a sequence, it can be de-constructed into n-mers to produce a set of internal contiguous subsequences. From any given target sequence, we would be able to determine what fragments would result. The hybridization sequence method depends, in part, upon being able to work in the reverse, from a set of fragments of known sequences to the full sequence. In simple cases, one is able to start at a single position and work in either or both directions towards the ends of the sequence as illustrated in the example.

The number of possible sequences of a given length increases very quickly with the length of that sequence. Thus, a 10-mer of zeros and ones has 1024 possibilities, a 12-mer has 4096. A 20-mer has over a million possibilities, and a 30-mer has over a billion. However, a given 30-mer has, at most, 26 different internal 5-mer sequences. Thus, a 30 character target sequence having over a million possible sequences can be substantially defined by only 26 different 5-mers. It will be recognized that the probe oligonucleotides will preferably, but need not necessarily, be of identical length, and that the probe sequences need not necessarily be contiguous in that the overlapping subsequences need not differ by only a single subunit. Moreover, each position of the matrix pattern need not be homogeneous, but may actually contain a plurality of probes of known sequence. In addition, although all of the possible subsequence specifications would be preferred, a less than full set of sequences specifications could be used. In particular, although a substantial fraction will preferably be at least about 70%, it may be less than that. About 20% would be preferred, more preferably at least about 30% would be preferred. Higher percentages would be especially preferred.



US 6,355,432 B1

15

## 2. Example of Four Letter Alphabet

A four letter alphabet may be conceptualized in at least two different ways from the two letter alphabet. One way is to consider the four possible values at each position and to analogize in a similar fashion to the binary example each of the overlaps. A second way is to group the binary digits into groups.

Using the first means, the overlap comparisons are performed with a four letter alphabet rather than a two letter alphabet. Then, in contrast to the binary system with 10 positions where  $2^{10}=1024$  possible sequences, in a 4-character alphabet with 10 positions, there will actually be  $4^{10}=1,048,576$  possible sequences. Thus, the complexity of a four character sequence has a much larger number of possible sequences compared to a two character sequence. Note, however, that there are still only 6 different internal 5-mers. For simplicity, we shall examine a 5 character string with 3 character subsequences. Instead of only 1 and 0, the characters may be designated, e.g., A, C, G, and T. Let us take the sequence GGCTA. The 3-mer subsequences are:

---

GGC  
GCT  
CTA

---

Given these subsequences, there is one sequence, or at most only a few sequences which would produce that combination of subsequences, i.e., GGCTA.

Alternatively, with a four character universe, the binary system can be looked at in pairs of digits. The pairs would be 00, 01, 10, and 11. In this manner, the earlier used sequence 1010011100 is looked at as 10,10,01,11,00. Then the first character of two digits is selected from the possible universe of the four representations 00, 01, 10, and 11. Then a probe would be in an even number of digits, e.g., not five digits, but, three pairs of digits or six digits. A similar comparison is performed and the possible overlaps determined. The 3-pair subsequences are:

10, 10, 01  
10, 01, 11  
01, 11, 00

and the overlap reconstruction produces 10,10,01,11,00.

The latter of the two conceptual views of the 4 letter alphabet provides a representation which is similar to what would be provided in a digital computer. The applicability to a four nucleotide alphabet is easily seen by assigning, e.g., 00 to A, 01 to C, 10 to G, and 11 to T. And, in fact, if such a correspondence is used, both examples for the 4 character sequences can be seen to represent the same target sequence. The applicability of the hybridization method and its analysis for determining the ultimate sequence is easily seen if A is the representation of adenine, C is the representation of cytosine, G is the representation of guanine, and T is the representation of thymine or uracil.

## 3. Generalization to m-letter Alphabet

This reconstruction process may be applied to polymers of virtually any number of possible characters in the alphabet, and for virtually any length sequence to be sequenced, though limitations, as discussed below, will limit its efficiency at various extremes of length. It will be recognized that the theory can be applied to a large diversity of systems where sequence is important.

16

For example, the method could be applied to sequencing of a polypeptide. A polypeptide can have any of twenty natural amino acid possibilities at each position. A twenty letter alphabet is amenable to sequencing by this method so long as reagents exist for recognizing shorter subsequences therein. A preferred reagent for achieving that goal would be a set of monoclonal antibodies each of which recognizes a specific three contiguous amino acid subsequence. A complete set of antibodies which recognize all possible subsequences of a given length, e.g., 3 amino acids, and preferably with a uniform affinity, would be  $20^3=8000$  reagents.

It will also be recognized that each target sequence which is recognized by the specific reagents need not have homogeneous termini. Thus, fragments of the entire target sequence will also be useful for hybridizing appropriate subsequences. It is, however, preferable that there not be a significant amount of labeled homogeneous contaminating extraneous sequences. This constraint does usually require the purification of the target molecule to be sequenced, but a specific label technique would dispense with a purification requirement if the unlabeled extraneous sequences do not interfere with the labeled sequences.

In addition, conformational effects of target polypeptide folding may, in certain embodiments, be negligible if the polypeptide is fragmented into sufficiently small peptides, or if the interaction is performed under conditions where conformation, but not specific interaction, is disrupted.

## B. Complications

Two obvious complications exist with the method of sequence analysis by hybridization. The first results from a probe of inappropriate length while the second relates to internally repeated sequences.

The first obvious complication is a problem which arises from an inappropriate length of recognition sequence, which causes problems with the specificity of recognition. For example, if the recognized sequence is too short, every sequence which is utilized will be recognized by every probe sequence. This occurs, e.g., in a binary system where the probes are each of sequences which occur relatively frequently, e.g., a two character probe for the binary system. Each possible two character probe would be expected to appear  $1/4$  of the time in every single two character position. Thus, the above sequence example would be recognized by each of the 00, 10, 01, and 11. Thus, the sequence information is virtually lost because the resolution is too low and each recognition reagent specifically binds at multiple sites on the target sequence.

The number of different probes which bind to a target depends on the relationship between the probe length and the target length. At the extreme of short probe length, the just mentioned problem exists of excessive redundancy and lack of resolution. The lack of stability in recognition will also be a problem with extremely short probes. At the extreme of long probe length, each entire probe sequence is on a different position of a substrate. However, a problem arises from the number of possible sequences, which goes up dramatically with the length of the sequence. Also, the specificity of recognition begins to decrease as the contribution to binding by any particular subunit may become sufficiently low that the system fails to distinguish the fidelity of recognition. Mismatched hybridization may be a problem with the polynucleotide sequencing applications, though the fingerprinting and mapping applications may not be so strict in their fidelity requirements. As indicated above, a thirty position binary sequence has over a million possible sequences, a number which starts to become unreasonably large in its required number of different sequences, even

US 6,355,432 B1

17

though the target length is still very short. Preparing a substrate with all sequence possibilities for a long target may be extremely difficult due to the many different oligomers which must be synthesized.

The above example illustrates how a long target sequence may be reconstructed with a reasonably small number of shorter subsequences. Since the present day resolution of the regions of the substrate having defined oligomer probes attached to the substrate approaches about 10 microns by 10 microns for resolvable regions, about  $10^6$ , or 1 million, positions can be placed on a one centimeter square substrate. However, high resolution systems may have particular disadvantages which may be outweighed using the lower density substrate matrix pattern. For this reason, a sufficiently large number of probe sequences can be utilized so that any given target sequence may be determined by hybridization to a relatively small number of probes.

A second complication relates to convergence of sequences to a single subsequence. This will occur when a particular subsequence is repeated in the target sequence. This problem can be addressed in at least two different ways. The first, and simpler way, is to separate the repeat sequences onto two different targets. Thus, each single target will not have the repeated sequence and can be analyzed to its end. This solution, however, complicates the analysis by requiring that some means for cutting at a site between the repeats can be located. Typically a careful sequencer would want to have two intermediate cut points so that the intermediate region can also be sequenced in both directions across each of the cut points. This problem is inherent in the hybridization method for sequencing but can be minimized by using a longer known probe sequence so that the frequency of probe repeats is decreased.

Knowing the sequence of flanking sequences of the repeat will simplify the use of polymerase chain reaction (PCR) or a similar technique to further definitively determine the sequence between sequence repeats. Probes can be made to hybridize to those known sequences adjacent the repeat sequences, thereby producing new target sequences for analysis. See, e.g., Innis et al. (eds.) (1990) *PCR Protocols: A Guide to Methods and Applications*, Academic Press; and methods for synthesis of oligonucleotide probes, see, e.g., Gait (1984) *Oligonucleotide Synthesis: A Practical Approach*, IRL Press, Oxford.

Other means for dealing with convergence problems include using particular longer probes, and using degeneracy reducing analogues, see, e.g., Macevitz, S. (1990) PCT publication number WO 90/04652, which is hereby incorporated herein by reference. By use of stretches of the degeneracy reducing analogues with other probes in particular combinations, the number of probes necessary to fully saturate the possible oligomer probes is decreased. For example, with a stretch of 12-mers having the central 4-mer of degenerate nucleotides, in combination with all of the possible 8-mers, the collection numbers twice the number of possible 8-mers, e.g.  $65,536 + 65,536 = 131,072$ , but the population provides screening equivalent to all possible 12-mers.

By way of further explanation, all possible oligonucleotide 8-mers may be depicted in the fashion:

N1-N2-N3-N4-N5-N6-N7-N8,

in which there are  $4^8 = 65,536$  possible 8-mers. As described in Ser. No. 07/624,120, now abandoned, producing all possible 8-mers requires  $4 \times 8 = 32$  chemical binary synthesis steps to produce the entire matrix pattern of 65,536 8-mer possibilities. By incorporating degeneracy reducing

18

nucleotides, D's, which hybridize nonselectively to any corresponding complementary nucleotide, new oligonucleotides 12-mers can be made in the fashion:

N1-N2-N3-N4-D-D-D-D-N5-N6-N7-N8,

in which there are again, as above, only  $4^8 = 65,536$  possible "12-mers", which in reality only have 8 different nucleotides.

However, it can be seen that each possible 12-mer probe could be represented by a group of the two 8-mer types. Moreover, repeats of less than 12 nucleotides would not converge, or cause repeat problems in the analysis. Thus, instead of requiring a collection of probes corresponding to all 12-mers, or  $4^{12} = 16,777,216$  different 12-mers, the same information can be derived by making 2 sets of "8-mers" consisting of the typical 8-mer collection of  $4^8 = 65,536$  and the "12-mer" set with the degeneracy reducing analogues, also requiring making  $4^8 = 65,536$ . The combination of the two sets, requires making  $65,536 + 65,536 = 131,072$  different molecules, but giving the information of 16,777,216 molecules. Thus, incorporating the degeneracy reducing analogue decreases the number of molecules necessary to get 12-mer resolution by a factor of about 128-fold.

#### C. Non-polynucleotide Embodiments

The above example is directed towards a polynucleotide embodiment. This application is relatively easily achieved because the specific reagents will typically be complementary oligonucleotides, although in certain embodiments other specific reagents may be desired. For example, there may be circumstances where other than complementary base pairing will be utilized. The polynucleotide targets, will usually be single strand, but may be double or triple stranded in various applications. However, a triple stranded specific interaction might be sometimes desired, or a protein or other specific binding molecule may be utilized. For example, various promoter or DNA sequence specific binding proteins might be used, including, e.g., restriction enzyme binding domains, other binding domains, and antibodies. Thus, specific recognition reagents besides oligonucleotides may be utilized.

For other polymer targets, the specific reagents will often be polypeptides. These polypeptides may be protein binding domains from enzymes or other proteins which display specificity for binding. Usually an antibody molecule may be used, and monoclonal antibodies may be particularly desired. Classical methods may be applied for preparing antibodies, see, e.g., Harlow and Lane (1988) *Antibodies: A Laboratory Manual* Cold Spring Harbor Press, New York; and Goding (1986) *Monoclonal Antibodies: Principles and Practice* (2d Ed.) Academic Press, San Diego. Other suitable techniques for in vitro exposure of lymphocytes to the antigens or selection of libraries of antibody binding sites are described, e.g., in Huse et al. (1989) *Science* 246:1275-1281; and Ward et al. (1989) *Nature* 341:544-546, each of which is hereby incorporated herein by reference. Unusual antibody production methods are also described, e.g., in Hendricks et al. (1989) *BioTechnology* 7:1271-1274; and Hiatt et al. (1989) *Nature* 342:76-78, each of which is hereby incorporated herein by reference. Other molecules which may exhibit specific binding interaction may be useful for attachment to a VLSIPS substrate by various methods, including the caged biotin methods, see, e.g., Ser. No. 07/435,316, now abandoned, and Barrett et al. (1993) U.S. Pat. No. 5,252,743.

The antibody specific reagents should be particularly useful for the polypeptide, carbohydrate, and synthetic poly-



US 6,355,432 B1

19

mer applications. Individual specific reagents might be generated by an automated process to generate the number of reagents necessary to advantageously use the high density positional matrix pattern. In an alternative approach, a plurality of hybridoma cells may be screened for their ability to bind to a VLSIPS matrix possessing the desired sequences whose binding specificity is desired. Each cell might be individually grown up and its binding specificity determined by VLSIPS apparatus and technology. An alternative strategy would be to expose the same VLSIPS matrix to a polyclonal serum of high titer. By a successively large volume of serum and different animals, each region of the VLSIPS substrate would have attached to it a substantial number of antibody molecules with specificity of binding. The substrate, with non-covalently bound antibodies could be derivatized and the antibodies transferred to an adjacent second substrate in the matrix pattern in which the antibody molecules had attached to the first matrix. If the sensitivity of detection of binding interaction is sufficiently high, such a low efficiency transfer of antibody molecules may produce a sufficiently high signal to be useful for many purposes, including the sequencing applications.

In another embodiment, capillary forces may be used to transfer the selected reagents to a new matrix, to which the reagents would be positionally attached in the pattern of the recognized sequences. Or, the reagents could be transversely electrophoresed, magnetically transferred, or otherwise transported to a new substrate in their retained positional pattern.

### III. POLYNUCLEOTIDE SEQUENCING

In principle, the making of a substrate having a positionally defined matrix pattern of all possible oligonucleotides of a given length involves a conceptually simple method of synthesizing each and every different possible oligonucleotide, and affixing them to a definable position. Oligonucleotide synthesis is presently mechanized and enabled by current technology, see, e.g., Ser. No. 07/362,901, now abandoned; Pirrung et al. (1992) U.S. Pat. No. 5,143,854; and instruments supplied by Applied Biosystems, Foster City, Calif.

#### A. Preparation of Substrate Matrix

The production of the collection of specific oligonucleotides used in polynucleotide sequencing may be produced in at least two different ways. Present technology certainly allows production of ten nucleotide oligomers on a solid phase or other synthesizing system. See, e.g., instrumentation provided by Applied Biosystems, Foster City, Calif. Although a single oligonucleotide can be relatively easily made, a large collection of them would typically require a fairly large amount of time and investment. For example, there are  $4^{10}=1,048,576$  possible ten nucleotide oligomers. Present technology allows making each and every one of them in a separate purified form though such might be costly and laborious.

Once the desired repertoire of possible oligomer sequences of a given length have been synthesized, this collection of reagents may be individually positionally attached to a substrate, thereby allowing a batchwise hybridization step. Present technology also would allow the possibility of attaching each and every one of these 10-mers to a separate specific position on a solid matrix. This attachment could be automated in any of a number of ways, particularly through the use of a caged biotin type linking. This would produce a matrix having each of different possible 10-mers.

A batchwise hybridization is much preferred because of its reproducibility and simplicity. An automated process of

20

attaching various reagents to positionally defined sites on a substrate is provided in Pirrung et al. (1992) U.S. Pat. No. 5,143,854; Ser. No. 07/624,120, now abandoned; and Barrett et al. (1993) U.S. Pat. No. 5,252,743; each of which is hereby incorporated herein by reference.

Instead of separate synthesis of each oligonucleotide, these oligonucleotides are conveniently synthesized in parallel by sequential synthetic processes on a defined matrix pattern as provided in Pirrung et al. (1992) U.S. Pat. No. 5,143,854; and Ser. No. 07/624,120, now abandoned, which are incorporated herein by reference. Here, the oligonucleotides are synthesized stepwise on a substrate at positionally separate and defined positions. Use of photosensitive blocking reagents allows for defined sequences of synthetic steps over the surface of a matrix pattern. By use of the binary masking strategy, the surface of the substrate can be positioned to generate a desired pattern of regions, each having a defined sequence oligonucleotide synthesized and immobilized thereto.

Although the prior art technology can be used to generate the desired repertoire of oligonucleotide probes, an efficient and cost effective means would be to use the VLSIPS technology described in Pirrung et al. (1992) U.S. Pat. No. 5,143,854 and Ser. No. 07/624,120, now abandoned. In this embodiment, the photosensitive reagents involved in the production of such a matrix are described below.

The regions for synthesis may be very small, usually less than about  $100\text{ }\mu\text{m}\times 100\text{ }\mu\text{m}$ , more usually less than about  $50\text{ }\mu\text{m}\times 50\text{ }\mu\text{m}$ . The photolithography technology allows synthetic regions of less than about  $10\text{ }\mu\text{m}\times 10\text{ }\mu\text{m}$ , about  $3\text{ }\mu\text{m}\times 3\text{ }\mu\text{m}$ , or less. The detection also may detect such sized regions, though larger areas are more easily and reliably measured.

At a size of about 30 microns by 30 microns, one million regions would take about 11 centimeters square or a single wafer of about 4 centimeters by 4 centimeters. Thus the present technology provides for making a single matrix of that size having all one million plus possible oligonucleotides. Region size is sufficiently small to correspond to densities of at least about 5 regions/cm<sup>2</sup>, 20 regions/cm<sup>2</sup>, 50 regions/cm<sup>2</sup>, 100 regions/cm<sup>2</sup>, and greater, including 300 regions/cm<sup>2</sup>, 1000 regions/cm<sup>2</sup>, 3K regions/cm<sup>2</sup>, 10K regions/cm<sup>2</sup>, 30K regions/cm<sup>2</sup>, 100K regions/cm<sup>2</sup>, 300K regions/cm<sup>2</sup> or more, even in excess of one million regions/cm<sup>2</sup>.

Although the pattern of the regions which contain specific sequences is theoretically not important, for practical reasons certain patterns will be preferred in synthesizing the oligonucleotides. The application of binary masking algorithms for generating the pattern of known oligonucleotide probes is described in related Ser. No. 07/624,120, now abandoned, which was filed simultaneously with this application. By use of these binary masks, a highly efficient means is provided for producing the substrate with the desired matrix pattern of different sequences. Although the binary masking strategy allows for the synthesis of all lengths of polymers, the strategy may be easily modified to provide only polymers of a given length. This is achieved by omitting steps where a subunit is not attached.

The strategy for generating a specific pattern may take any of a number of different approaches. These approaches are well described in related application Ser. No. 07/624,120, now abandoned, and include a number of binary masking approaches which will not be exhaustively discussed herein. However, the binary masking and binary synthesis approaches provide a maximum of diversity with a minimum number of actual synthetic steps.

US 6,355,432 B1

21

The length of oligonucleotides used in sequencing applications will be selected on criteria determined to some extent by the practical limits discussed above. For example, if probes are made as oligonucleotides, there will be 65,536 possible eight nucleotide sequences. If a nine subunit oligonucleotide is selected, there are 262,144 possible permutations of sequences. If a ten-mer oligonucleotide is selected, there are 1,048,576 possible permutations of sequences. As the number gets larger, the required number of positionally defined subunits necessary to saturate the possibilities also increases. With respect to hybridization conditions, the length of the matching necessary to confer stability of the conditions selected can be compensated for. See, e.g., Kanehisa, M. (1984) *Nuc. Acids Res.* 12:203-213, which is hereby incorporated herein by reference.

Although not described in detail here, but below for oligonucleotide probes, the VLSIPS technology would typically use a photosensitive protective group on an oligonucleotide. Sample oligonucleotides are shown in FIG. 1. In particular, the photoprotective group on the nucleotide molecules may be selected from a wide variety of positive light reactive groups preferably including nitro aromatic compounds such as o-nitro-benzyl derivatives or benzylsulfonyl. See, e.g., Gait (1984) *Oligonucleotide Synthesis: A Practical Approach*, IRL Press, Oxford, which is hereby incorporated herein by reference. In a preferred embodiment, 6-nitro-veratryl oxycarbonyl (NVOC), 2-nitrobenzyl oxycarbonyl (NBOC), or  $\alpha,\alpha$ -dimethyl-dimethoxybenzyl oxycarbonyl (DEZ) is used. Photoremovable protective groups are described in, e.g., Patchornik (1970) *J. Amer. Chem. Soc.* 92:6333-6335; and Amit et al. (1974) *J. Organic Chem.* 39:192-196; each of which is hereby incorporated herein by reference.

A preferred linker for attaching the oligonucleotide to a silicon matrix is illustrated in FIG. 2. A more detailed description is provided below. A photosensitive blocked nucleotide may be attached to specific locations of unblocked prior cycles of attachments on the substrate and can be successively built up to the correct length oligonucleotide probe.

It should be noted that multiple substrates may be simultaneously exposed to a single target sequence where each substrate is a duplicate of one another or where, in combination, multiple substrates together provide the complete or desired subset of possible subsequences. This provides the opportunity to overcome a limitation of the density of positions on a single substrate by using multiple substrates. In the extreme case, each probe might be attached to a single bead or substrate and the beads sorted by whether there is a binding interaction. Those beads which do bind might be encoded to indicate the subsequence specificity of reagents attached thereto.

Then, the target may be bound to the whole collection of beads and those beads that have appropriate specific reagents on them will bind to the target. Then a sorting system may be utilized to sort those beads that actually bind the target from those that do not. This may be accomplished by presently available cell sorting devices or a similar apparatus. After the relatively small number of beads which have bound the target have been collected, the encoding scheme may be read off to determine the specificity of the reagent on the bead. An encoding system may include, a magnetic system, a shape encoding system, a color encoding system, or a combination of any of these, or any other encoding system. Once again, with the collection of specific interactions that have occurred, the binding may be analyzed for sequence information, fingerprint information, or mapping information.

22

The parameters of polynucleotide sizes of both the probes and target sequences are determined by the applications and other circumstances. The length of the oligonucleotide probes used will depend in part upon the limitations of the VLSIPS technology to provide the number of desired probes. For example, in an absolute sequencing application, it is often useful to have virtually all of the possible oligonucleotides of a given length. As indicated above, there are 65,536 8-mers, 262,144 9-mers, 1,048,576 10-mers, 4,194,304 11-mers, etc. As the length of the oligomer increases the number of different probes which must be synthesized also increases at a rate of a factor of 4 for every additional nucleotide. Eventually the size of the matrix and the limitations in the resolution of regions in the matrix will reach the point where an increase in number of probes becomes disadvantageous. However, this sequencing procedure requires that the system be able to distinguish, by appropriate selection of hybridization and washing conditions, between binding of absolute fidelity and binding of complementary sequences containing mismatches. On the other hand, if the fidelity is unnecessary, this discrimination is also unnecessary and a significantly longer probe may be used. Significantly longer probes would typically be useful in fingerprinting or mapping applications.

The length of the probe is selected for a length that will allow the probe to bind with specificity to possible targets. The hybridization conditions are also very important in that they will determine how closely the homology of complementary binding will be detected. In fact, a single target may be evaluated at a number of different conditions to determine its spectrum of specificity for binding particular probes. This may find use in a number of other applications besides the polynucleotide sequencing fingerprinting or mapping. For example, it will be desired to determine the spectrum of binding affinities and specificities of cell surface antigens with binding by particular antibodies immobilized on the substrate surface, particularly under different interaction conditions. In a related fashion, different regions with reagents having differing affinities or levels of specificity may allow such a spectrum to be defined using a single incubation, where various regions, at a given hybridization condition, show the binding affinity. For example, fingerprint probes of various lengths, or with specific defined non-matches may be used. Unnatural nucleotides or nucleotides exhibiting modified specificity of complementary binding are described in greater detail in Macevitz (1990) PCT pub. No. WO 90/04652; and see the section on modified nucleotides in the Sigma Chemical Company catalogue.

#### B. Labeling Target Nucleotide

The label used to detect the target sequences will be determined, in part, by the detection methods being applied. Thus, the labeling method and label used are selected in combination with the actual detecting systems being used. Once a particular label has been selected, appropriate labeling protocols will be applied, as described below for specific embodiments. Standard labeling protocols for nucleic acids are described, e.g., in Sambrook et al.; Kambara, H. et al. (1988) *BioTechnology* 6:816-821; Smith, L. et al. (1985) *Nuc. Acids Res.* 13:2399-2412; for polypeptides, see, e.g., Allen G. (1989) *Sequencing of Proteins and Peptides*, Elsevier, New York, especially chapter 5, and Greenstein and Winitz (1961) *Chemistry of the Amino Acids*, Wiley and Sons, New York. Carbohydrate labeling is described, e.g., in Chaplin and Kennedy (1986) *Carbohydrate Analysis: A Practical Approach*, IRL Press, Oxford. Labeling of other polymers will be performed by methods applicable to them as recognized by a person having ordinary skill in manipulating the corresponding polymer.

US 6,355,432 B1

23

In some embodiments, the target need not actually be labeled if a means for detecting where interaction takes place is available. As described below, for a nucleic acid embodiment, such may be provided by an intercalating dye which intercalates only into double stranded segments, e.g., where interaction occurs. See, e.g., Sheldon et al. U.S. Pat. No. 4,582,789.

In many uses, the target sequence will be absolutely homogeneous, both with respect to the total sequence and with respect to the ends of each molecule. Homogeneity with respect to sequence is important to avoid ambiguity. It is preferable that the target sequences of interest not be contaminated with a significant amount of labeled contaminating sequences. The extent of allowable contamination will depend on the sensitivity of the detection system and the inherent signal to noise of the system. Homogeneous contamination sequences will be particularly disruptive of the sequencing procedure.

However, although the target polynucleotide must have a unique sequence, the target molecules need not have identical ends. In fact, the homogeneous target molecule preparation may be randomly sheared to increase the numerical number of molecules. Since the total information content remains the same, the shearing results only in a higher number of distinct sequences which may be labeled and bind to the probe. This fragmentation may give a vastly superior signal relative to a preparation of the target molecules having homogeneous ends.

The signal for the hybridization is likely to be dependent on the numerical frequency of the target-probe interactions. If a sequence is individually found on a larger number of separate molecules a better signal will result. In fact, shearing a homogeneous preparation of the target may often be preferred before the labeling procedure is performed, thereby producing a large number of labeling groups associated with each subsequence.

#### C. Hybridization Conditions

The hybridization conditions between probe and target should be selected such that the specific recognition interaction, i.e., hybridization, of the two molecules is both sufficiently specific and sufficiently stable. See, e.g., Hames and Higgins (1985) *Nucleic Acid Hybridisation: A Practical Approach*, IRL Press, Oxford. These conditions will be dependent both on the specific sequence and often on the guanine and cytosine (GC) content of the complementary hybrid strands. The conditions may often be selected to be universally equally stable independent of the specific sequences involved. This typically will make use of a reagent such as an alkylammonium buffer. See, Wood et al. (1985) "Base Composition-independent Hybridization in Tetramethylammonium Chloride: A Method for Oligonucleotide Screening of Highly Complex Gene Libraries," *Proc. Natl. Acad. Sci. USA*, 82:1585-1588; and Krupov et al. (1989) "An Oligonucleotide Hybridization Approach to DNA Sequencing," *FEBS Letters*, 256:118-122; each of which is hereby incorporated herein by reference. An alkylammonium buffer tends to minimize differences in hybridization rate and stability due to GC content. By virtue of the fact that sequences then hybridize with approximately equal affinity and stability, there is relatively little bias in strength or kinetics of binding for particular sequences. Temperature and salt conditions along with other buffer parameters should be selected such that the kinetics of renaturation should be essentially independent of the specific target subsequence or oligonucleotide probe involved. In order to ensure this, the hybridization reactions will usually be performed in a single incubation of all the substrate matrices

24

together exposed to the identical same target probe solution under the same conditions.

Alternatively, various substrates may be individually treated differently. Different substrates may be produced, each having reagents which bind to target subsequences with substantially identical stabilities and kinetics of hybridization. For example, all of the high GC content probes could be synthesized on a single substrate which is treated accordingly. In this embodiment, the arylammonium buffers could be unnecessary. Each substrate is then treated in a manner such that the collection of substrates show essentially uniform binding and the hybridization data of target binding to the individual substrate matrix is combined with the data from other substrates to derive the necessary subsequence binding information. The hybridization conditions will usually be selected to be sufficiently specific such that the fidelity of base matching will be properly discriminated. Of course, control hybridizations should be included to determine the stringency and kinetics of hybridization.

#### D. Detection; VLSIPS™ Technology Scanning

The next step of the sequencing process by hybridization involves labeling of target polynucleotide molecules. A quickly and easily detectable signal is preferred. The VLSIPS™ Technology apparatus is designed to easily detect a fluorescent label, so fluorescent tagging of the target sequence is preferred. Other suitable labels include heavy metal labels, magnetic probes, chromogenic labels (e.g., phosphorescent labels, dyes, and fluorophores) spectroscopic labels, enzyme linked labels, radioactive labels, and labeled binding proteins. Additional labels are described in U.S. Pat. No. 4,366,241, which is incorporated herein by reference.

The detection methods used to determine where hybridization has taken place will typically depend upon the label selected above. Thus, for a fluorescent label a fluorescent detection step will typically be used. Pirrung et al. (1992) U.S. Pat. No. 5,143,854 and Ser. No. 07/624,120, now abandoned, describe apparatus and mechanisms for scanning a substrate matrix using fluorescence detection, but a similar apparatus is adaptable for other optically detectable labels.

The detection method provides a positional localization of the region where hybridization has taken place. However, the position is correlated with the specific sequence of the probe since the probe has specifically been attached or synthesized at a defined substrate matrix position. Having collected all of the data indicating the subsequences present in the target sequence, this data may be aligned by overlap to reconstruct the entire sequence of the target, as illustrated above.

It is also possible to dispense with actual labeling if some means for detecting the positions of interaction between the sequence specific reagent and the target molecule are available. This may take the form of an additional reagent which can indicate the sites either of interaction, or the sites of lack of interaction, e.g., a negative label. For the nucleic acid embodiments, locations of double strand interaction may be detected by the incorporation of intercalating dyes, or other reagents such as antibody or other reagents that recognize helix formation, see, e.g., Sheldon, et al. (1986) U.S. Pat. No. 4,582,789, which is hereby incorporated herein by reference.

#### E. Analysis

Although the reconstruction can be performed manually as illustrated above, a computer program will typically be used to perform the overlap analysis. A program may be written and run on any of a large number of different



US 6,355,432 B1

25

computer hardware systems. The variety of operating systems and languages useable will be recognized by a computer software engineer. Various different languages may be used, e.g., BASIC; C; PASCAL; etc. A simple flow chart of data analysis is illustrated in FIG. 1.

#### F. Substrate Reuse

Finally, after a particular sequence has been hybridized and the pattern of hybridization analyzed, the matrix substrate should be reusable and readily prepared for exposure to a second or subsequent target polynucleotides. In order to do so, the hybrid duplexes are disrupted and the matrix treated in a way which removes all traces of the original target. The matrix may be treated with various detergents or solvents to which the substrate, the oligonucleotide probes, and the linkages to the substrate are inert. This treatment may include an elevated temperature treatment, treatment with organic or inorganic solvents, modifications in pH, and other means for disrupting specific interaction. Thereafter, a second target may actually be applied to the recycled matrix and analyzed as before.

#### G. Non-Polynucleotide Aspects

Although the sequencing, fingerprinting, and mapping functions will make use of the natural sequence recognition property of complementary nucleotide sequences, the non-polynucleotide sequences typically require other sequence recognition reagents. These reagents will take the form, typically, of proteins exhibiting binding specificity, e.g., enzyme binding sites or antibody binding sites.

Enzyme binding sites may be derived from promoter proteins, restriction enzymes, and the like. See, e.g., Stryer, L. (1988) *Biochemistry*, W.H. Freeman, Palo Alto. Antibodies will typically be produced using standard procedures, see, e.g., Harlow and Lane (1988) *Antibodies: A Laboratory Manual*, Cold Spring Harbor Press, New York; and Goding (1986) *Monoclonal Antibodies: Principles and Practice*, (2d Ed.) Academic Press, San Diego.

Typically, an antigen, or collection of antigens are presented to an immune system. This may take the form of synthesized short polymers produced by the VLSIPS technology, or by the other synthetic means, or from isolation of natural products. For example, antigen for the polypeptides may be made by the VLSIPS technology, by standard peptide synthesis, by isolation of natural proteins with or without degradation to shorter segments, or by expression of a collection of short nucleic acids of random or defined sequences. See, e.g., Tuerk and Gold (1990) *Science* 249:505-510, for generation of a collection of randomly mutagenized oligonucleotides useful for expression.

The antigen or collection is presented to an appropriate immune system, e.g., to a whole animal as in a standard immunization protocol, or to a collection of immune cells or equivalent. In particular, see Ward et al. (1989) *Nature* 341:544-546; and Huse et al. (1989) *Science* 246:1275-1281, each of which is hereby incorporated herein by reference.

A large diversity of antibodies will be generated, some of which have specificities for the desired sequences. Antibodies may be purified having the desired sequence specificities by isolating the cells producing them. For example, a VLSIPS substrate with the desired antigens synthesized thereon may be used to isolate cells with cell surface reagents which recognize the antigens. The VLSIPS substrate may be used as an affinity reagent to select and recover the appropriate cells. Antibodies from those cells may be attached to a substrate using the caged biotin methodology, or by attaching a targeting molecule, e.g., an oligonucle-

26

otide. Alternatively, the supernatants from antibody producing cells can be easily assayed using a VLSIPS substrate to identify the cells producing the appropriate antibodies.

Although cells may be isolated, specific antibody molecules which perform the sequence recognition will also be sufficient. Preferably populations of antibody with a known specificity can be isolated. Supernatants from a large population of producing cells may be passed over a VLSIPS substrate to bind to the desired antigens attached to the substrate. When a sufficient density of antibody molecules are attached, they may be removed by an automated process, preferably as antibody populations exhibiting specificity of binding.

In one particular embodiment, a VLSIPS substrate, e.g., with a large plurality of fingerprint antigens attached thereto, is used to isolate antibodies from a supernatant of a population of cells producing antibodies to the antigens. Using the substrate as an affinity reagent, the antibodies will attach to the appropriate positionally defined antigens. The antibodies may be carefully removed therefrom, preferably by an automated system which retains their homogeneous specificities. The isolated antibodies can be attached to a new substrate in a positionally defined matrix pattern.

In a further embodiment, these spatially separated antibodies may be isolated using a specific targeting method for isolation. In this embodiment, a linker molecule which attaches to a particular portion of the antibody, preferably away from the binding site, can be attached to the antibodies. Various reagents will be used, including staphylococcus protein A or antibodies which bind to domains remote from the binding site. Alternatively, the antibodies in the population, before affinity purification, may be derivatized with an appropriate reagent compatible with new VLSIPS synthesis. A preferred reagent is a nucleotide which can serve as a linker to synthetic VLSIPS steps for synthesizing a specific sequence thereon. Then, by successive VLSIPS cycles, each of the antibodies attached to the defined antigen regions can have a defined oligonucleotide synthesized thereon and corresponding in area to the region of the substrate having each antigen attached. These defined oligonucleotides will be useful as targeting reagents to attach those antibodies possessing the same target sequence specificity at defined positions on a new substrate, by virtue of having bound to the antigen region, to a new VLSIPS substrate having the complementary target oligonucleotides positionally located on it. In this fashion, a VLSIPS substrate having the desired antigens attached thereto can be used to generate a second VLSIPS substrate with positionally defined reagents which recognize those antigens.

The selected antigens will typically be selected to be those which define particular functionalities or properties, so as to be useful for fingerprinting and other uses. They will also be useful for mapping and sequencing embodiments.

#### IV. FINGERPRINTING

##### A. General

Many of the procedures and techniques used in the polynucleotide sequencing section are also appropriate for fingerprinting applications. See, e.g., Poustka, et al. (1986) *Cold Spring Harbor Symposia on Quant. Biol.*, vol. LI, 131-139, Cold Spring Harbor Press, New York; which is hereby incorporated herein by reference. The fingerprinting method provided herein is based, in part, upon the ability to positionally localize a large number of different specific probes onto a single substrate. This high density matrix pattern provides the ability to screen for, or detect, a very large number of different sequences simultaneously. In fact, depending upon the hybridization conditions, fingerprinting

US 6,355,432 B1

27

to the resolution of virtually absolute matching of sequence is possible thereby approaching an absolute sequencing embodiment. And the sequencing embodiment is very useful in identifying the probes useful in further fingerprinting uses. For example, characteristic features of genetic sequences will be identified as being diagnostic of the entire sequence. However, in most embodiments, longer probe and target will be used, and for which slight mismatching may not need to be resolved.

#### B. Preparation of Substrate Matrix

A collection of specific probes may be produced by either of the methods described above in the section on sequencing. Specific oligonucleotide probes of desired lengths may be individually synthesized on a standard oligonucleotide synthesizer. The length of these probes is limited only by the ability of the synthesizer to continue to accurately synthesize a molecule. Oligonucleotides or sequence fragments may also be isolated from natural sources. Biological amplification methods may be coupled with synthetic synthesizing procedures such as, e.g., polymerase chain reaction.

In one embodiment, the individually isolated probes may be attached to the matrix at defined positions. These probe reagents may be attached by an automated process making use of the caged biotin methodology described in Ser. No. 07/612,671, or using photochemical reagents, see, e.g., Dattagupta et al. (1985) U.S. Pat. No. 4,542,102 and (1987) U.S. Pat. No. 4,713,326. Each individually purified reagent can be attached individually at specific locations on a substrate.

In another embodiment, the VLSIPS synthesizing technique may be used to synthesize the desired probes at specific positions on a substrate. The probes may be synthesized by successively adding appropriate monomer subunits, e.g., nucleotides, to generate the desired sequences.

In another embodiment, a relatively short specific oligonucleotide is used which serves as a targeting reagent for positionally directing the sequence recognition reagent. For example, the sequence specific reagents having a separate additional sequence recognition segment (usually of a different polymer from the target sequence) can be directed to target oligonucleotides attached to the substrate. By use of non-natural targeting reagents, e.g., unusual nucleotide analogues which pair with other unnatural nucleotide analogues and which do not interfere with natural nucleotide interactions, the natural and non-natural portions can coexist on the same molecule without interfering with their individual functionalities. This can combine both a synthetic and biological production system analogous to the technique for targeting monoclonal antibodies to locations on a VLSIPS substrate at defined positions. Unnatural optical isomers of nucleotides may be useful unnatural reagents subject to similar chemistry, but incapable of interfering with the natural biological polymers. See also, Ser. No. 07/626,730, which is hereby incorporated herein by reference.

After the separate substrate attached reagents are attached to the targeting segment, the two are crosslinked, thereby permanently attaching them to the substrate. Suitable crosslinking reagents are known, see, e.g., Dattagupta et al. (1985) U.S. Pat. No. 4,542,102 and (1987) "Coupling of nucleic acids to solid support by photochemical methods," U.S. Pat. No. 4,713,326, each of which is hereby incorporated herein by reference. Similar linkages for attachment of proteins to a solid substrate are provided, e.g., in Merrifield (1986) *Science* 232:341-347, which is hereby incorporated herein by reference.

28

#### C. Labeling Target Nucleotides

The labeling procedures used in the sequencing embodiments will also be applicable in the fingerprinting embodiments. However, since the fingerprinting embodiments often will involve relatively large target molecules and relatively short oligonucleotide probes, the amount of signal necessary to incorporate into the target sequence may be less critical than in the sequencing applications. For example, a relatively long target with a relatively small number of labels per molecule may be easily amplified or detected because of the relatively large target molecule size.

In various embodiments, it may be desired to cleave the target into smaller segments as in the sequencing embodiments. The labeling procedures and cleavage techniques described in the sequencing embodiments would usually also be applicable here.

#### D. Hybridization Conditions

The hybridization conditions used in fingerprinting embodiments will typically be less critical than for the sequencing embodiments. The reason is that the amount of mismatching which may be useful in providing the fingerprinting information would typically be far greater than that necessary in sequencing uses. For example, Southern hybridizations do not typically distinguish between slightly mismatched sequences. Under these circumstances, important and valuable information may be arrived at with less stringent hybridization conditions while providing valuable fingerprinting information. However, since the entire substrate is typically exposed to the target molecule at one time, the binding affinity of the probes should usually be of approximately comparable levels. For this reason, if oligonucleotide probes are being used, their lengths should be approximately comparable and will be selected to hybridize under conditions which are common for most of the probes on the substrate. Much as in a Southern hybridization, the target and oligonucleotide probes are of lengths typically greater than about 25 nucleotides. Under appropriate hybridization conditions, e.g., typically higher salt and lower temperature, the probes will hybridize irrespective of imperfect complementarity. In fact, with probes of greater than, e.g., about fifty nucleotides, the difference in stability of different sized probes will be relatively minor.

Typically the fingerprinting is merely for probing similarity or homology. Thus, the stringency of hybridization can usually be decreased to fairly low levels. See, e.g., Wetmur and Davidson (1968) "Kinetics of Renaturation of DNA," *J. Mol. Biol.*, 31:349-370; and Kanehisa, M. (1984) *Nuc. Acids Res.*, 12:203-213.

#### E. Detection; VLSIPS™ Technology Scanning

Detection methods will be selected which are appropriate for the selected label. The scanning device need not necessarily be digitized or placed into a specific digital database, though such would most likely be done. For example, the analysis in fingerprinting could be photographic. Where a standardized fingerprint substrate matrix is used, the pattern of hybridizations may be spatially unique and may be compared photographically. In this manner, each sample may have a characteristic pattern of interactions and the likelihood of identical patterns will preferably be such low frequency that the fingerprint pattern indeed becomes a characteristic pattern virtually as unique as an individual's fingertip fingerprint. With a standardized substrate, every individual could be, in theory, uniquely identifiable on the basis of the pattern of hybridizing to the substrate.

Of course, the VLSIPS™ Technology scanning apparatus may also be useful to generate a digitized version of the fingerprint pattern. In this way, the identification pattern can